

# Today...

- Approximations and the shapes of graphs.
- Hill functions.
- Motivating limits: secant lines, tangent lines.

# Which of the following is a safe approximation to make?

- (A) If  $a$  is small then we can say  $a \approx 0$ .
- (B) If  $a$  is small then we can say  $ab \approx 0$ .
- (C) If  $a$  is small then we can say  $a+b \approx b$ .
- (D) If  $a$  is small compared to  $b$  then we can say  $a+b \approx b$ .

# Which of the following is a safe approximation to make?

- (A) If  $a$  is small then we can say  $a \approx 0$ .
- (B) If  $a$  is small then we can say  $ab \approx 0$ .
- (C) If  $a$  is small then we can say  $a+b \approx b$ .
- (D) If  $a$  is small compared to  $b$  then we can say  $a+b \approx b$ .

Explain notation  $a \ll b$ .

# Which of the following is a safe approximation to make?

- (A) If  $a$  is small then we can say  $a \approx 0$ .
- (B) If  $a$  is small then we can say  $ab \approx 0$ .
- (C) If  $a$  is small then we can say  $a+b \approx b$ .
- (D) If  $a$  is small compared to  $b$  then we can say  $a+b \approx b$ .

# Which of the following is a safe approximation to make?

- (A) If  $a$  is small then we can say  $a \approx 0$ .
- (B) If  $a$  is small then we can say  $ab \approx 0$ .
- (C) If  $a$  is small then we can say  $a+b \approx b$ .
- (D) If  $a$  is small compared to  $b$  then we can say  $a+b \approx b$ .

Explain notation  $a \ll b$ .

# Comparisons and approximation must be based on relative sizes!

For each of the following, (A) True, (B) False . . . You line up some bricks to make a wall one brick high.

- One brick is a small number of bricks (i.e. “a is small”).

# Comparisons and approximation must be based on relative sizes!

For each of the following, (A) True, (B) False . . . You line up some bricks to make a wall one brick high.

- One brick is a small number of bricks (i.e. “a is small”).
- The wall is small ( $a \approx 0$ ).

# Comparisons and approximation must be based on relative sizes!

For each of the following, (A) True, (B) False . . . You line up some bricks to make a wall one brick high.

- One brick is a small number of bricks (i.e. “a is small”).
- The wall is small ( $a \approx 0$ ).
- If you make the wall 20 times as high, it is still small ( $ab \approx 0$ ).



# Comparisons and approximation must be based on relative sizes!

For each of the following, (A) True, (B) False . . . You line up some bricks to make a wall one brick high.

- One brick is a small number of bricks (i.e. “a is small”).
- The wall is small ( $a \approx 0$ ).
- If you make the wall 20 times as high, it is still small ( $ab \approx 0$ ).
- Add one row of bricks to a wall 20 bricks high. The new wall about the same size as the old wall. ( $a+b \approx b$ )

# Comparisons and approximation must be based on relative sizes!

For each of the following, (A) True, (B) False . . . You line up some bricks to make a wall one brick high.

- One brick is a small number of bricks (i.e. “a is small”). **True.**
- The wall is small ( $a \approx 0$ ).
- If you make the wall 20 times as high, it is still small ( $ab \approx 0$ ).
- Add one row of bricks to a wall 20 bricks high. The new wall about the same size as the old wall. ( $a+b \approx b$ )

# Comparisons and approximation must be based on relative sizes!

For each of the following, (A) True, (B) False . . . You line up some bricks to make a wall one brick high.

- One brick is a small number of bricks (i.e. “a is small”). **True.**
- The wall is small ( $a \approx 0$ ). **Depends.**
- If you make the wall 20 times as high, it is still small ( $ab \approx 0$ ).
- Add one row of bricks to a wall 20 bricks high. The new wall about the same size as the old wall. ( $a+b \approx b$ )

# Comparisons and approximation must be based on relative sizes!

For each of the following, (A) True, (B) False . . . You line up some bricks to make a wall one brick high.

- One brick is a small number of bricks (i.e. “a is small”). **True.**
- The wall is small ( $a \approx 0$ ). **Depends.**
- If you make the wall 20 times as high, it is still small ( $ab \approx 0$ ). **Depends.**
- Add one row of bricks to a wall 20 bricks high. The new wall about the same size as the old wall. ( $a+b \approx b$ )

# Comparisons and approximation must be based on relative sizes!

For each of the following, (A) True, (B) False . . . You line up some bricks to make a wall one brick high.

- One brick is a small number of bricks (i.e. “a is small”). **True.**
- The wall is small ( $a \approx 0$ ). **Depends.**
- If you make the wall 20 times as high, it is still small ( $ab \approx 0$ ). **Depends.**
- Add one row of bricks to a wall 20 bricks high. The new wall about the same size as the old wall. ( $a+b \approx b$ ) **True.**



# Comparisons and approximation must be based on relative sizes!

- For
- son
- On
- The
- If y
- Ad
- ab



[http://commons.wikimedia.org/wiki/File:Big\\_foot.JPG](http://commons.wikimedia.org/wiki/File:Big_foot.JPG)

up  
ue.  
0).  
ends.  
all



# Comparisons and approximation must be based on relative sizes!

For each of the following, (A) True, (B) False . . . You line up some bricks to make a wall one brick high.

- One brick is a small number of bricks (i.e. “a is small”). **True.**
- The wall is small ( $a \approx 0$ ). **Depends.**
- If you make the wall 20 times as high, it is still small ( $ab \approx 0$ ). **Depends.**
- Add one row of bricks to a wall 20 bricks high. The new wall about the same size as the old wall. ( $a+b \approx b$ ) **True.**

# Comparisons and approximation must be based on relative sizes!

For  
son

- On
- The
- If y
- Ad
- ab



[http://commons.wikimedia.org/wiki/File:Talia\\_on\\_her\\_pink\\_Strider.jpg](http://commons.wikimedia.org/wiki/File:Talia_on_her_pink_Strider.jpg)

up

ue.

0).  
ends.

all



**When  $0 < x \ll b$ , then  $x + b$  can be approximated by...**

(A)  $b$

(B)  $x$

(C) infinity

**When  $0 < x \ll b$ , then  $x + b$  can be approximated by...**

(A)  $b$

(B)  $x$

(C) infinity

**When  $0 < x \ll b$ , then  $f(x) = \frac{ax^n}{b^n + x^n}$**

**can be approximated by...**

(A)  $a$

(B)  $\frac{a}{b^n}$

(C)  $a \left(\frac{x}{b}\right)^n$

(D)  $0$

(E)  $1$

**When  $0 < x \ll b$ , then  $f(x) = \frac{ax^n}{b^n + x^n}$**

**can be approximated by...**

(A)  $a$

(B)  $\frac{a}{b^n}$

(C)  $a \left(\frac{x}{b}\right)^n$

(D)  $0$

(E)  $1$

**When  $x \gg b$ , then  $x + b$  can be approximated by...**

(A)  $b$

(B)  $x$

(C) infinity

**When  $x \gg b$ , then  $x + b$  can be approximated by...**

(A)  $b$

(B)  $x$

(C) infinity

**When  $x \gg b$ , then  $f(x) = \frac{ax^n}{b^n + x^n}$**

**can be approximated by...**

(A)  $a$

(B)  $\frac{a}{b^n}$

(C)  $a \left(\frac{x}{b}\right)^n$

(D)  $0$

(E)  $1$

**When  $x \gg b$ , then  $f(x) = \frac{ax^n}{b^n + x^n}$**

**can be approximated by...**

(A)  $a$

(B)  $\frac{a}{b^n}$

(C)  $a \left(\frac{x}{b}\right)^n$


(D)  $0$

(E)  $1$




# Implications for graphing

$$f(x) = \frac{ax^n}{b^n + x^n}$$




# Implications for graphing

$$f(x) = \frac{ax^n}{b^n + x^n}$$




# Implications for graphing

$$f(x) = \frac{ax^n}{b^n + x^n}$$




# Implications for graphing

$$f(x) = \frac{ax^n}{b^n + x^n}$$




# Implications for graphing

$$f(x) = \frac{ax^n}{b^n + x^n}$$




# Implications for graphing

$$f(x) = \frac{ax^n}{b^n + x^n}$$




# Implications for graphing

$$f(x) = \frac{ax^n}{b^n + x^n}$$



# Implications for graphing


$$f(x) = \frac{ax^n}{b^n + x^n}$$





# Implications for graphing

$$f(x) = \frac{ax^n}{b^n + x^n}$$



# Comparing Hill functions with different n values


(A) Green:  $n=2$ , yellow:  $n=3$ , red:  $n=4$ , blue:  $n=5$ .

(B) Green:  $n=4$ , yellow:  $n=3$ , red:  $n=2$ , blue:  $n=1$ .

(C) Green:  $n=5$ , yellow:  $n=4$ , red:  $n=3$ , blue:  $n=2$ .

(D) Either (B) or (C) (not enough info).

$$f(x) = \frac{ax^n}{b^n + x^n}$$



# Comparing Hill functions with different n values


(A) Green:  $n=2$ , yellow:  $n=3$ , red:  $n=4$ , blue:  $n=5$ .

(B) Green:  $n=4$ , yellow:  $n=3$ , red:  $n=2$ , blue:  $n=1$ .

(C) Green:  $n=5$ , yellow:  $n=4$ , red:  $n=3$ , blue:  $n=2$ .

(D) Either (B) or (C) (not enough info).

$$f(x) = \frac{ax^n}{b^n + x^n}$$



**What is the slope of the line connecting the points?**

(A)  $m = (x_1 - x_2) / (y_1 - y_2)$

•  $(x_2, y_2)$

(B)  $m = (x_2 - x_1) / (y_1 - y_2)$

(C)  $m = (y_1 - y_2) / (x_1 - x_2)$

•  
 $(x_1, y_1)$

(D)  $m = (y_2 - y_1) / (x_2 - x_1)$

**What is the slope of the line connecting the points?**

(A)  $m = (x_1 - x_2) / (y_1 - y_2)$

•  $(x_2, y_2)$

(B)  $m = (x_2 - x_1) / (y_1 - y_2)$

(C)  $m = (y_1 - y_2) / (x_1 - x_2)$

•  $(x_1, y_1)$

(D)  $m = (y_2 - y_1) / (x_2 - x_1)$


# What is the slope of the secant line to the graph of $f(x)$ ?

(A)  $m = (f(x_1) - f(x_2)) / (x_2 - x_1)$

(B)  $m = (f(x_2) - f(x_1)) / (x_2 - x_1)$

(C)  $m = (x_1 - x_2) / (f(x_1) - f(x_2))$

(D)  $m = (x_2 - x_1) / (f(x_1) - f(x_2))$



Slope of secant line = **average rate of change** from  $x_1$  to  $x_2$ .


# What is the slope of the secant line to the graph of $f(x)$ ?

(A)  $m = (f(x_1) - f(x_2)) / (x_2 - x_1)$

(B)  $m = (f(x_2) - f(x_1)) / (x_2 - x_1)$


(C)  $m = (x_1 - x_2) / (f(x_1) - f(x_2))$

(D)  $m = (x_2 - x_1) / (f(x_1) - f(x_2))$



Slope of secant line = **average rate of change** from  $x_1$  to  $x_2$ .


**What if you want the rate of change AT  $x_1$ ?**






# What if you want the rate of change AT $x_1$ ?

Take a point  $x_2$  so that the secant line is closer to the “secant line” AT  $x_1$ .




# What if you want the rate of change AT $x_1$ ?

Take a point  $x_2$  so that the secant line is closer to the “secant line” AT  $x_1$ .




# What if you want the rate of change AT $x_1$ ?

Take a point  $x_2$  so that the secant line is closer to the “secant line” AT  $x_1$ .




# What if you want the rate of change AT $x_1$ ?

Take a point  $x_2$  so that the secant line is closer to the “secant line” AT  $x_1$ .



# What if you want the rate of change AT $x_1$ ?

Take a point  $x_2$  so that the secant line is closer to the “secant line” AT  $x_1$ .




Alternate notation: let  $x_2 = x_1 + h$  so that

$$m = \frac{f(x_2) - f(x_1)}{x_2 - x_1}$$

# What if you want the rate of change AT $x_1$ ?

Take a point  $x_2$  so that the secant line is closer to the “secant line” AT  $x_1$ .




Alternate notation: let  $x_2 = x_1 + h$  so that

$$m = \frac{f(x_2) - f(x_1)}{x_2 - x_1} = \frac{f(x_1 + h) - f(x_1)}{x_1 + h - x_1}$$

# What if you want the rate of change AT $x_1$ ?

Take a point  $x_2$  so that the secant line is closer to the “secant line” AT  $x_1$ .



Alternate notation: let  $x_2 = x_1 + h$  so that

$$m = \frac{f(x_2) - f(x_1)}{x_2 - x_1} = \frac{f(x_1 + h) - f(x_1)}{h}$$

# If we take $h$ values closer and closer to 0...

- The secant line approaches the **tangent line**.
- The slope of the secant line approaches the slope of the tangent line.
- We call the resulting slope **the derivative at  $x_1$** .
- We now have to learn how to take **limits!**

$$\text{slope at } x_1 = f'(x_1) = \lim_{h \rightarrow 0} \frac{f(x_1 + h) - f(x_1)}{h}$$