

Today

- Foraging – a complicated example that I'll use to emphasize the value of thinking about the problem in biological, graphical and formulaic terms.
- Reminders:
 - Assignment 5 due Thursday
 - No class on Monday (Thanksgiving)
 - OSH 4 due Wednesday
 - Regular PLQs.

Foraging


Foraging time includes

- a commute ($t_0 \rightarrow$ constant),
- a visit to each patch (nt_p)



Foraging

Foraging success is characterized by $f(t_p)$ = resource extracted from a single patch after a time t_p spent in the patch.



Remember the definition of $f(t_p)$ for an upcoming clicker Q.

Foraging

- (A) When food is scarce, it is best to maximize $R(t)$.
- (B) When food is scarce, it is best to maximize $E(t)$.
- (C) When food is abundant, with many competing priorities to deal with, it is best to maximize $E(t)$.
- (D) Maximizing $E(t)$ and maximizing $R(t)$ are the same.
- (E) When food is abundant, it is best to maximize $E(t)$.





Foraging

- (A) When food is scarce, it is best to maximize $R(t)$.
- (B) When food is scarce, it is best to maximize $E(t)$.
- (C) When food is abundant, with many competing priorities to deal with, it is best to maximize $E(t)$.
- (D) Maximizing $E(t)$ and maximizing $R(t)$ are the same.
- (E) When food is abundant, it is best to maximize $E(t)$.



Foraging

Maximize average rate of resource extraction, that is, how fast the squirrel gathers food.




If the squirrel visits n patches, each for t_p minutes, total time spent foraging is...

(A) $t_{\text{tot}} = nt_p$

(B) $t_{\text{tot}} = nt_0$

(C) $t_{\text{tot}} = nt_0 + t_p$

(D) $t_{\text{tot}} = nt_p + t_0$




If the squirrel visits n patches, each for t_p minutes, total time spent foraging is...

(A) $t_{\text{tot}} = nt_p$

(B) $t_{\text{tot}} = nt_0$

(C) $t_{\text{tot}} = nt_0 + t_p$

(D) $t_{\text{tot}} = nt_p + t_0$




If the squirrel visits n patches,
each for t_p minutes, total
resource extracted is...

(A) $r = nf(t_p)$

(B) $r = f(nt_p)$

(C) $r = f(nt_p + t_0)$

(D) $r = nf(t_p + t_0)$



Foraging

Foraging success is characterized by $f(t_p)$ = resource extracted from a single patch after a time t_p spent in the patch.



Remember the definition of $f(t_p)$ for an upcoming clicker Q.


If the squirrel visits n patches,
each for t_p minutes, total
resource extracted is...

(A) $r = nf(t_p)$

(B) $r = f(nt_p)$

(C) $r = f(nt_p + t_0)$

(D) $r = nf(t_p + t_0)$




If the squirrel visits n patches,
each for t_p minutes, total
resource extracted is...

(A) $r = nf(t_p)$

(B) $r = f(nt_p)$


(C) $r = f(nt_p + t_0)$

(D) $r = nf(t_p + t_0)$








Average rate of resource extraction if t_p is spent in each patch:

$$R_{\text{avg}}(t_p) = \text{total extracted} / \text{total time}$$
$$= nf(t_p) / (nt_p + t_0)$$








Which of the following graphs matches the given description of $f(t)$?

Collection goes well at first but gradually slows down as the resource is depleted.








Which of the following graphs matches the given description of $f(t)$?

Collection goes well at first but gradually slows down as the resource is depleted.








Which of the following graphs matches the given description of $f(t)$?

It is initially hard to find nuts but gets easier with time. Eventually, there are none left to collect.








Which of the following graphs matches the given description of $f(t)$?

It is initially hard to find nuts but gets easier with time. Eventually, there are none left to collect.








Which of the following graphs matches the given description of $f(t)$?

Initially, you collect some nuts but the birds figure out what you're doing and start stealing from you.




Which of the following graphs matches the given description of $f(t)$?

Initially, you collect some nuts but the birds figure out what you're doing and start stealing from you.



Choose $f(t_p) = \text{constant} = C$

Find t_p that maximizes $R_{\text{avg}} = nC / (nt_p + t_0)$



(A) $t_p = -t_0/n$


(B) $t_p = 0$

(C) Never leave.

Think and/or sketch before you calculate.

Choose $f(t_p) = \text{constant} = C$

Find t_p that maximizes $R_{\text{avg}} = nC / (nt_p + t_0)$



(A) $t_p = -t_0/n$

(B) $t_p = 0$


(C) Never leave.

Think and/or sketch before you calculate.

Choose $f(t_p) = t_p^2$

Find t_p that maximizes $R_{avg} = nt_p^2 / (nt_p + t_0)$

"maximum"



(A) $t_p = 2nt_0$


(B) $t_p = 0$

(C) Never leave.

Think and/or sketch before you calculate.

Choose $f(t_p) = t_p^2$

Find t_p that maximizes $R_{avg} = nt_p^2 / (nt_p + t_0)$




Think and/or sketch before you calculate.

Choose $f(t_p) = t_p / (k + t_p)$

Find t_p that maximizes

$$R_{\text{avg}} = nt_p / (nt_p + t_0) (k + t_p)$$



(A) $t_p = 0$

(B) $t_p = \text{sqrt}(kt_0/n)$


(C) Never leave.

Think and/or sketch before you calculate.

Choose $f(t_p) = t_p / (k + t_p)$

Find t_p that maximizes

$$R_{avg} = nt_p / (nt_p + t_0) (k + t_p)$$



(A) $t_p = 0$

(B) $t_p = \text{sqrt}(kt_0/n)$

(C) Never leave.

Think and/or sketch before you calculate.

Summary message:

- It is useful to think about the physical problem and the function being optimized before jumping into calculating derivatives and finding absolute extrema.