


Today

- Modeling with delta-function forcing (tanks, springs)
- Convolution
- Transfer functions


Delta-function forcing (6.5)

- Water with $c_{in} = 2$ g/L of sugar enters a tank at a rate of $r = 1$ L/min. The initially sugar-free tank holds $V = 5$ L and the contents are well-mixed. Water drains from the tank at a rate r . At $t_{cube} = 3$ min, a sugar cube of mass $m_{cube} = 3$ g is dropped into the tank.




Delta-function forcing (6.5)

- Water with $c_{in} = 2$ g/L of sugar enters a tank at a rate of $r = 1$ L/min. The initially sugar-free tank holds $V = 5$ L and the contents are well-mixed. Water drains from the tank at a rate r . At $t_{cube} = 3$ min, a sugar cube of mass $m_{cube} = 3$ g is dropped into the tank.
 - Sketch the mass of salt in the tank as a function of time (from intuition).



Delta-function forcing (6.5)


- Water with $c_{in} = 2$ g/L of sugar enters a tank at a rate of $r = 1$ L/min. The initially sugar-free tank holds $V = 5$ L and the contents are well-mixed. Water drains from the tank at a rate r . At $t_{cube} = 3$ min, a sugar cube of mass $m_{cube} = 3$ g is dropped into the tank.
 - Sketch the mass of salt in the tank as a function of time (from intuition).
 - Write down an ODE for the mass of sugar in the tank as a function of time.



Delta-function forcing (6.5)

- Water with $c_{in} = 2$ g/L of sugar enters a tank at a rate of $r = 1$ L/min. The initially sugar-free tank holds $V = 5$ L and the contents are well-mixed. Water drains from the tank at a rate r . At $t_{cube} = 3$ min, a sugar cube of mass $m_{cube} = 3$ g is dropped into the tank.
 - Sketch the mass of salt in the tank as a function of time (from intuition).
 - Write down an ODE for the mass of sugar in the tank as a function of time.


$$m' = r c_{in} - \frac{r}{V} m + m_{cube} \delta(t - t_{cube})$$



Delta-function forcing (6.5)

- Water with $c_{in} = 2$ g/L of sugar enters a tank at a rate of $r = 1$ L/min. The initially sugar-free tank holds $V = 5$ L and the contents are well-mixed. Water drains from the tank at a rate r . At $t_{cube} = 3$ min, a sugar cube of mass $m_{cube} = 3$ g is dropped into the tank.
 - Sketch the mass of salt in the tank as a function of time (from intuition).
 - Write down an ODE for the mass of sugar in the tank as a function of time.

$$m' = r c_{in} - \frac{r}{V} m + m_{cube} \delta(t - t_{cube})$$




- Note: $\delta(t)$ has units of 1/time.

Delta-function forcing (6.5)

- Water with $c_{in} = 2$ g/L of sugar enters a tank at a rate of $r = 1$ L/min. The initially sugar-free tank holds $V = 5$ L and the contents are well-mixed. Water drains from the tank at a rate r . At $t_{cube} = 3$ min, a sugar cube of mass $m_{cube} = 3$ g is dropped into the tank.
 - Sketch the mass of salt in the tank as a function of time (from intuition).
 - Write down an ODE for the mass of sugar in the tank as a function of time.

$$m' = r c_{in} - \frac{r}{V} m + m_{cube} \delta(t - t_{cube})$$

$$m' = 2 - \frac{1}{5} m + 3 \delta(t - 3)$$




Delta-function forcing (6.5)

- Water with $c_{in} = 2$ g/L of sugar enters a tank at a rate of $r = 1$ L/min. The initially sugar-free tank holds $V = 5$ L and the contents are well-mixed. Water drains from the tank at a rate r . At $t_{cube} = 3$ min, a sugar cube of mass $m_{cube} = 3$ g is dropped into the tank.
 - Sketch the mass of salt in the tank as a function of time (from intuition).
 - Write down an ODE for the mass of sugar in the tank as a function of time.

$$m' = r c_{in} - \frac{r}{V} m + m_{cube} \delta(t - t_{cube})$$

$$m' = 2 - \frac{1}{5} m + 3 \delta(t - 3)$$

- Solve the ODE.



Delta-function forcing (6.5)


- Water with $c_{in} = 2$ g/L of sugar enters a tank at a rate of $r = 1$ L/min. The initially sugar-free tank holds $V = 5$ L and the contents are well-mixed. Water drains from the tank at a rate r . At $t_{cube} = 3$ min, a sugar cube of mass $m_{cube} = 3$ g is dropped into the tank.
 - Sketch the mass of salt in the tank as a function of time (from intuition).
 - Write down an ODE for the mass of sugar in the tank as a function of time.

$$m' = r c_{in} - \frac{r}{V} m + m_{cube} \delta(t - t_{cube})$$

$$m' = 2 - \frac{1}{5} m + 3 \delta(t - 3)$$

- Solve the ODE.

$$m(t) = 10(1 - e^{-t/5}) + 3u_3(t)e^{-(t-3)/5}$$



Delta-function forcing (6.5)

- Water with $c_{in} = 2$ g/L of sugar enters a tank at a rate of $r = 1$ L/min. The initially sugar-free tank holds $V = 5$ L and the contents are well-mixed. Water drains from the tank at a rate r . At $t_{cube} = 3$ min, a sugar cube of mass $m_{cube} = 3$ g is dropped into the tank.
 - Sketch the mass of salt in the tank as a function of time (from intuition).
 - Write down an ODE for the mass of sugar in the tank as a function of time.


$$m' = rc_{in} - \frac{r}{V}m + m_{cube}\delta(t - t_{cube})$$

$$m' = 2 - \frac{1}{5}m + 3\delta(t - 3)$$

- Solve the ODE.

$$m(t) = 10(1 - e^{-t/5}) + 3u_3(t)e^{-(t-3)/5}$$

- Sketch the solution to the ODE. How would it differ if $t_{cube}=10$ min?



Delta-function forcing (6.5)

- Sketch the solution to the ODE.

$$m(t) = 10(1 - e^{-t/5}) + 3u_3(t)e^{-(t-3)/5}$$

Delta-function forcing (6.5)


- Sketch the solution to the ODE.

$$\begin{aligned} m(t) &= 10(1 - e^{-t/5}) + 3u_3(t)e^{-(t-3)/5} \\ &= \begin{cases} 10(1 - e^{-t/5}) & \text{for } t < 3, \\ 10 - (10 - 3e^{3/5})e^{-t/5} & \text{for } t \geq 3. \end{cases} \end{aligned}$$

Delta-function forcing (6.5)

- Sketch the solution to the ODE.


$$\begin{aligned} m(t) &= 10(1 - e^{-t/5}) + 3u_3(t)e^{-(t-3)/5} \\ &= \begin{cases} 10(1 - e^{-t/5}) & \text{for } t < 3, \\ 10 - (10 - 3e^{3/5})e^{-t/5} & \text{for } t \geq 3. \end{cases} \end{aligned}$$



Delta-function forcing (6.5)

- Sketch the solution to the ODE.


$$\begin{aligned} m(t) &= 10(1 - e^{-t/5}) + 3u_3(t)e^{-(t-3)/5} \\ &= \begin{cases} 10(1 - e^{-t/5}) & \text{for } t < 3, \\ 10 - (10 - 3e^{3/5})e^{-t/5} & \text{for } t \geq 3. \end{cases} \end{aligned}$$



Delta-function forcing (6.5)

- Sketch the solution to the ODE.


$$\begin{aligned} m(t) &= 10(1 - e^{-t/5}) + 3u_3(t)e^{-(t-3)/5} \\ &= \begin{cases} 10(1 - e^{-t/5}) & \text{for } t < 3, \\ 10 - (10 - 3e^{3/5})e^{-t/5} & \text{for } t \geq 3. \end{cases} \end{aligned}$$



Delta-function forcing (6.5)

- Sketch the solution to the ODE.


$$m(t) = 10(1 - e^{-t/5}) + 3u_3(t)e^{-(t-3)/5}$$
$$= \begin{cases} 10(1 - e^{-t/5}) & \text{for } t < 3, \\ 10 - (10 - 3e^{3/5})e^{-t/5} & \text{for } t \geq 3. \end{cases}$$



Delta-function forcing (6.5)

- Sketch the solution to the ODE.


$$m(t) = 10(1 - e^{-t/5}) + 3u_3(t)e^{-(t-3)/5}$$
$$= \begin{cases} 10(1 - e^{-t/5}) & \text{for } t < 3, \\ 10 - (10 - 3e^{3/5})e^{-t/5} & \text{for } t \geq 3. \end{cases}$$



Delta-function forcing (6.5)

- Sketch the solution to the ODE.


$$\begin{aligned} m(t) &= 10(1 - e^{-t/5}) + 3u_3(t)e^{-(t-3)/5} \\ &= \begin{cases} 10(1 - e^{-t/5}) & \text{for } t < 3, \\ 10 - (10 - 3e^{3/5})e^{-t/5} & \text{for } t \geq 3. \end{cases} \end{aligned}$$



Delta-function forcing (6.5)

- Sketch the solution to the ODE.


$$m(t) = 10(1 - e^{-t/5}) + 3u_3(t)e^{-(t-3)/5}$$
$$= \begin{cases} 10(1 - e^{-t/5}) & \text{for } t < 3, \\ 10 - (10 - 3e^{3/5})e^{-t/5} & \text{for } t \geq 3. \end{cases}$$



Delta-function forcing (6.5)

- Sketch the solution to the ODE.


$$m(t) = 10(1 - e^{-t/5}) + 3u_3(t)e^{-(t-3)/5}$$
$$= \begin{cases} 10(1 - e^{-t/5}) & \text{for } t < 3, \\ 10 - (10 - 3e^{3/5})e^{-t/5} & \text{for } t \geq 3. \end{cases}$$



Delta-function forcing (6.5)

- Sketch the solution to the ODE.

$$\begin{aligned} m(t) &= 10(1 - e^{-t/5}) + 3u_3(t)e^{-(t-3)/5} \\ &= \begin{cases} 10(1 - e^{-t/5}) & \text{for } t < 3, \\ 10 - (10 - 3e^{3/5})e^{-t/5} & \text{for } t \geq 3. \end{cases} \end{aligned}$$




Delta-function forcing (6.5)

- Sketch the solution to the ODE.

$$m(t) = 10(1 - e^{-t/5}) + 3u_3(t)e^{-(t-3)/5}$$
$$= \begin{cases} 10(1 - e^{-t/5}) & \text{for } t < 3, \\ 10 - (10 - 3e^{3/5})e^{-t/5} & \text{for } t \geq 3. \end{cases}$$

- How would it differ if $t_{\text{cube}}=10$ min?




Delta-function forcing (6.5)

- Sketch the solution to the ODE.

$$m(t) = 10(1 - e^{-t/5}) + 3u_3(t)e^{-(t-3)/5}$$
$$= \begin{cases} 10(1 - e^{-t/5}) & \text{for } t < 3, \\ 10 - (10 - 3e^{3/5})e^{-t/5} & \text{for } t \geq 3. \end{cases}$$

- How would it differ if $t_{\text{cube}}=10$ min?




Delta-function forcing (6.5)

- Sketch the solution to the ODE.

$$m(t) = 10(1 - e^{-t/5}) + 3u_3(t)e^{-(t-3)/5}$$
$$= \begin{cases} 10(1 - e^{-t/5}) & \text{for } t < 3, \\ 10 - (10 - 3e^{3/5})e^{-t/5} & \text{for } t \geq 3. \end{cases}$$

- How would it differ if $t_{\text{cube}}=10$ min?




Delta-function forcing (6.5)

- Sketch the solution to the ODE.

$$m(t) = 10(1 - e^{-t/5}) + 3u_3(t)e^{-(t-3)/5}$$
$$= \begin{cases} 10(1 - e^{-t/5}) & \text{for } t < 3, \\ 10 - (10 - 3e^{3/5})e^{-t/5} & \text{for } t \geq 3. \end{cases}$$

- How would it differ if $t_{\text{cube}}=10$ min?




Delta-function forcing (6.5)

- Sketch the solution to the ODE.

$$m(t) = 10(1 - e^{-t/5}) + 3u_3(t)e^{-(t-3)/5}$$
$$= \begin{cases} 10(1 - e^{-t/5}) & \text{for } t < 3, \\ 10 - (10 - 3e^{3/5})e^{-t/5} & \text{for } t \geq 3. \end{cases}$$

- How would it differ if $t_{\text{cube}}=10$ min?




Delta-function forcing (6.5)

- Sketch the solution to the ODE.

$$m(t) = 10(1 - e^{-t/5}) + 3u_3(t)e^{-(t-3)/5}$$
$$= \begin{cases} 10(1 - e^{-t/5}) & \text{for } t < 3, \\ 10 - (10 - 3e^{3/5})e^{-t/5} & \text{for } t \geq 3. \end{cases}$$

- How would it differ if $t_{\text{cube}}=10$ min?




Delta-function forcing (6.5)

- A hammer hits a mass-spring system imparting an impulse of $I_0 = 2 \text{ N s}$ at $t = 5 \text{ s}$. The mass of the block is $m = 1 \text{ kg}$. The drag coefficient is $\gamma = 2 \text{ kg/s}$ and the spring constant is $k = 10 \text{ kg/s}^2$. The mass is initially at $y(0) = 2 \text{ m}$ with velocity $y'(0) = 0 \text{ m/s}$.


Delta-function forcing (6.5)

- A hammer hits a mass-spring system imparting an impulse of $I_0 = 2 \text{ N s}$ at $t = 5 \text{ s}$. The mass of the block is $m = 1 \text{ kg}$. The drag coefficient is $\gamma = 2 \text{ kg/s}$ and the spring constant is $k = 10 \text{ kg/s}^2$. The mass is initially at $y(0) = 2 \text{ m}$ with velocity $y'(0) = 0 \text{ m/s}$.



Delta-function forcing (6.5)

- A hammer hits a mass-spring system imparting an impulse of $I_0 = 2 \text{ N s}$ at $t = 5 \text{ s}$. The mass of the block is $m = 1 \text{ kg}$. The drag coefficient is $\gamma = 2 \text{ kg/s}$ and the spring constant is $k = 10 \text{ kg/s}^2$. The mass is initially at $y(0) = 2 \text{ m}$ with velocity $y'(0) = 0 \text{ m/s}$.
- Write down an equation for the position of the mass.



Delta-function forcing (6.5)


- A hammer hits a mass-spring system imparting an impulse of $I_0 = 2 \text{ N s}$ at $t = 5 \text{ s}$. The mass of the block is $m = 1 \text{ kg}$. The drag coefficient is $\gamma = 2 \text{ kg/s}$ and the spring constant is $k = 10 \text{ kg/s}^2$. The mass is initially at $y(0) = 2 \text{ m}$ with velocity $y'(0) = 0 \text{ m/s}$.
- Write down an equation for the position of the mass.

(A) $y'' + 2y' + 10y = 2 u_0(t)$

(B) $y'' + 2y' + 10y = 2 u_5(t)$

(C) $y'' + 2y' + 10y = 2 \delta(t)$

(D) $y'' + 2y' + 10y = 2 \delta(t - 5)$



Delta-function forcing (6.5)


- A hammer hits a mass-spring system imparting an impulse of $I_0 = 2 \text{ N s}$ at $t = 5 \text{ s}$. The mass of the block is $m = 1 \text{ kg}$. The drag coefficient is $\gamma = 2 \text{ kg/s}$ and the spring constant is $k = 10 \text{ kg/s}^2$. The mass is initially at $y(0) = 2 \text{ m}$ with velocity $y'(0) = 0 \text{ m/s}$.
- Write down an equation for the position of the mass.

(A) $y'' + 2y' + 10y = 2 u_0(t)$

(B) $y'' + 2y' + 10y = 2 u_5(t)$

(C) $y'' + 2y' + 10y = 2 \delta(t)$

★ (D) $y'' + 2y' + 10y = 2 \delta(t - 5)$



Delta-function forcing (6.5)

- A hammer hits a mass-spring system imparting an impulse of $I_0 = 2 \text{ N s}$ at $t = 5 \text{ s}$. The mass of the block is $m = 1 \text{ kg}$. The drag coefficient is $\gamma = 2 \text{ kg/s}$ and the spring constant is $k = 10 \text{ kg/s}^2$. The mass is initially at $y(0) = 2 \text{ m}$ with velocity $y'(0) = 0 \text{ m/s}$.
- Write down an equation for the position of the mass.


(A) $y'' + 2y' + 10y = 2 u_0(t)$

(B) $y'' + 2y' + 10y = 2 u_5(t)$

(C) $y'' + 2y' + 10y = 2 \delta(t)$

★ (D) $y'' + 2y' + 10y = 2 \delta(t - 5)$

$$s^2 Y - 2s + 2sY - 4 + 10Y = 2e^{-5c}$$



Delta-function forcing (6.5)

- A hammer hits a mass-spring system imparting an impulse of $I_0 = 2 \text{ N s}$ at $t = 5 \text{ s}$. The mass of the block is $m = 1 \text{ kg}$. The drag coefficient is $\gamma = 2 \text{ kg/s}$ and the spring constant is $k = 10 \text{ kg/s}^2$. The mass is initially at $y(0) = 2 \text{ m}$ with velocity $y'(0) = 0 \text{ m/s}$.
- Write down an equation for the position of the mass.

(A) $y'' + 2y' + 10y = 2 u_0(t)$


(B) $y'' + 2y' + 10y = 2 u_5(t)$

(C) $y'' + 2y' + 10y = 2 \delta(t)$

★ (D) $y'' + 2y' + 10y = 2 \delta(t - 5)$

$$s^2 Y - 2s + 2sY - 4 + 10Y = 2e^{-5s}$$

$$Y(s) = \frac{2(e^{-5s} + s + 2)}{s^2 + 2s + 10}$$



Delta-function forcing (6.5)

- Inverting $Y(s)$... (go through this on your own)

$$\begin{aligned} Y(s) &= \frac{2(e^{-5s} + s + 2)}{s^2 + 2s + 10} = \frac{2e^{-5s}}{s^2 + 2s + 10} + 2\frac{s + 2}{s^2 + 2s + 10} \\ &= \frac{2e^{-5s}}{s^2 + 2s + 10} + 2\frac{s + 2}{(s + 1)^2 + 9} \\ &= \frac{2e^{-5s}}{s^2 + 2s + 10} + 2\frac{s + 1}{(s + 1)^2 + 9} + \frac{2}{(s + 1)^2 + 9} \\ &= \frac{2e^{-5s}}{s^2 + 2s + 10} + 2\frac{s + 1}{(s + 1)^2 + 9} + \frac{2}{3}\frac{3}{(s + 1)^2 + 9} \\ &= \frac{2}{3}\frac{3e^{-5s}}{(s + 1)^2 + 9} + 2\frac{s + 1}{(s + 1)^2 + 9} + \frac{2}{3}\frac{3}{(s + 1)^2 + 9} \end{aligned}$$

$$y(t) = \frac{2}{3}u_5(t)e^{-(t-5)}\sin(3(t-5)) + 2e^{-t}\cos(3t) + \frac{2}{3}e^{-t}\sin(3t)$$

particular solution from δ forcing

homogeneous part

Convolution (6.6)

- We often end up with transforms to invert that are the product of two known transforms. For example,

$$Y(s) = \frac{2}{s^2(s^2 + 4)} = \frac{1}{s^2} \cdot \frac{2}{s^2 + 4}$$

Convolution (6.6)

- We often end up with transforms to invert that are the product of two known transforms. For example,

$$Y(s) = \frac{2}{s^2(s^2 + 4)} = \frac{1}{s^2} \cdot \frac{2}{s^2 + 4}$$

- Can we express the inverse of a product in terms of the known pieces?

$$F(s)G(s) = \mathcal{L}\{??\}$$

Convolution (6.6)

- We often end up with transforms to invert that are the product of two known transforms. For example,

$$Y(s) = \frac{2}{s^2(s^2 + 4)} = \frac{1}{s^2} \cdot \frac{2}{s^2 + 4}$$

- Can we express the inverse of a product in terms of the known pieces?

$$F(s)G(s) = \mathcal{L}\{??\}$$

$$F(s) = \int_0^{\infty} e^{-st} f(t) dt$$

Convolution (6.6)

- We often end up with transforms to invert that are the product of two known transforms. For example,

$$Y(s) = \frac{2}{s^2(s^2 + 4)} = \frac{1}{s^2} \cdot \frac{2}{s^2 + 4}$$

- Can we express the inverse of a product in terms of the known pieces?

$$F(s)G(s) = \mathcal{L}\{??\}$$

$$F(s) = \int_0^{\infty} e^{-st} f(t) dt$$

$$G(s) = \int_0^{\infty} e^{-st} g(t) dt$$

Convolution (6.6)

- We often end up with transforms to invert that are the product of two known transforms. For example,

$$Y(s) = \frac{2}{s^2(s^2 + 4)} = \frac{1}{s^2} \cdot \frac{2}{s^2 + 4}$$

- Can we express the inverse of a product in terms of the known pieces?

$$F(s)G(s) = \mathcal{L}\{??\}$$

$$F(s) = \int_0^{\infty} e^{-st} f(t) dt \quad \rightarrow \quad F(s) = \int_0^{\infty} e^{-s\tau} f(\tau) d\tau$$

$$G(s) = \int_0^{\infty} e^{-st} g(t) dt$$

Convolution (6.6)

- We often end up with transforms to invert that are the product of two known transforms. For example,

$$Y(s) = \frac{2}{s^2(s^2 + 4)} = \frac{1}{s^2} \cdot \frac{2}{s^2 + 4}$$

- Can we express the inverse of a product in terms of the known pieces?

$$F(s)G(s) = \mathcal{L}\{??\}$$

$$\begin{aligned} F(s) &= \int_0^{\infty} e^{-st} f(t) dt & \rightarrow & F(s) = \int_0^{\infty} e^{-s\tau} f(\tau) d\tau \\ G(s) &= \int_0^{\infty} e^{-st} g(t) dt & \rightarrow & G(s) = \int_0^{\infty} e^{-sw} g(w) dw \end{aligned}$$


Convolution (6.6)

$$F(s)G(s) = \int_0^{\infty} e^{-s\tau} f(\tau) d\tau \int_0^{\infty} e^{-sw} g(w) dw$$



Convolution (6.6)

$$F(s)G(s) = \int_0^{\infty} e^{-s\tau} f(\tau) d\tau \int_0^{\infty} e^{-sw} g(w) dw$$


$$= \int_0^{\infty} e^{-sw} g(w) \int_0^{\infty} e^{-s\tau} f(\tau) d\tau dw$$

$$= \int_0^{\infty} g(w) \int_0^{\infty} e^{-s(\tau+w)} f(\tau) d\tau dw$$

Replace τ using $u = \tau + w$ where w is constant in the inner integral.

$$= \int_0^{\infty} g(w) \int_w^{\infty} e^{-s(u)} f(u - w) du dw$$

$$= \int_0^{\infty} \int_w^{\infty} e^{-su} g(w) f(u - w) du dw$$


$$= \int_a^b \int_c^d e^{-su} g(w) f(u - w) dw du$$

Convolution (6.6)

- What are the correct values for a, b, c and d?

$$\int_0^{\infty} \int_w^{\infty} e^{-su} g(w) f(u-w) du dw$$

$$= \int_a^b \int_c^d e^{-su} g(w) f(u-w) dw du$$




Convolution (6.6)

- What are the correct values for a, b, c and d?

$$\int_0^{\infty} \int_w^{\infty} e^{-su} g(w) f(u-w) du dw$$

w=constant

$$= \int_a^b \int_c^d e^{-su} g(w) f(u-w) dw du$$




Convolution (6.6)

- What are the correct values for a, b, c and d?

$$\int_0^\infty \int_w^\infty e^{-su} g(w) f(u-w) du dw$$

(Note: In the original image, blue annotations are present: a blue arrow points from the inner integral's upper limit ∞ to the label $u=\infty$; another blue arrow points from the inner integral's lower limit w to the label $u=w$; a blue bracket under the entire double integral is labeled $w=\text{constant}$.)

$$= \int_a^b \int_c^d e^{-su} g(w) f(u-w) dw du$$




Convolution (6.6)

- What are the correct values for a, b, c and d?

$$\int_0^{\infty} \int_w^{\infty} e^{-su} g(w) f(u-w) du dw$$

(Note: In the original image, blue arrows point from the upper limit ∞ of the inner integral to the label $u=\infty$, and from the lower limit w to the label $u=w$. A blue bracket under the inner integral is labeled $w=\text{constant}$.)

$$= \int_a^b \int_c^d e^{-su} g(w) f(u-w) dw du$$




Convolution (6.6)

- What are the correct values for a, b, c and d?

$$\int_0^\infty \int_w^\infty e^{-su} g(w) f(u-w) du dw$$

(Note: In the original image, blue arrows point from the upper limit ∞ of the inner integral to the label $u=\infty$, and from the lower limit w to the label $u=w$. A blue bracket under the inner integral is labeled $w=\text{constant}$.)

$$= \int_a^b \int_c^d e^{-su} g(w) f(u-w) dw du$$




Convolution (6.6)

- What are the correct values for a, b, c and d?

$$\int_0^{\infty} \int_w^{\infty} e^{-su} g(w) f(u-w) du dw$$

(Note: In the original image, blue arrows point from the upper limit ∞ of the inner integral to the label $u=\infty$, and from the lower limit w to the label $u=w$. A blue arc under the inner integral is labeled $w=\text{constant}$.)

$$= \int_a^b \int_c^d e^{-su} g(w) f(u-w) dw du$$




Convolution (6.6)

- What are the correct values for a, b, c and d?

$$\int_0^{\infty} \int_w^{\infty} e^{-su} g(w) f(u-w) du dw$$

(Note: In the original image, blue annotations are present: 'u=' above the upper limit of the inner integral, 'u=w' below the lower limit of the inner integral, and a blue arrow labeled 'w=constant' pointing from the inner integral to the outer integral.)

$$= \int_a^b \int_c^d e^{-su} g(w) f(u-w) dw du$$




Convolution (6.6)

- What are the correct values for a, b, c and d?

$$\int_0^{\infty} \int_w^{\infty} e^{-su} g(w) f(u-w) du dw$$

$\underbrace{\hspace{15em}}_{w=\text{constant}}$

$$= \int_a^b \int_c^d e^{-su} g(w) f(u-w) dw du$$



- (A) Integrate in u from a=0 to b=∞ and in w from c=u, d=∞.
- (B) Integrate in u from a=0 to b=w and in w from c=0 to d=∞.
- (C) Integrate in u from a=0 to b=∞ and in w from c=0 to d=u.
- (D) Integrate in u from a=0 to b=∞ and in w from c=w to d=∞.
- (E) Huh?


Convolution (6.6)

- What are the correct values for a, b, c and d?

$$\int_0^{\infty} \int_w^{\infty} e^{-su} g(w) f(u-w) du dw$$

w=constant

$$= \int_a^b \int_c^d e^{-su} g(w) f(u-w) dw du$$




- (A) Integrate in u from a=0 to b=∞ and in w from c=u, d=∞.
- (B) Integrate in u from a=0 to b=w and in w from c=0 to d=∞.
- ★ (C) Integrate in u from a=0 to b=∞ and in w from c=0 to d=u.
- (D) Integrate in u from a=0 to b=∞ and in w from c=w to d=∞.
- (E) Huh?

Convolution (6.6)

- What are the correct values for a, b, c and d?

$$\int_0^{\infty} \int_w^{\infty} e^{-su} g(w) f(u-w) du dw$$




$$= \int_a^b \int_c^d e^{-su} g(w) f(u-w) dw du$$

- (A) Integrate in u from a=0 to b=∞ and in w from c=u, d=∞.
- (B) Integrate in u from a=0 to b=w and in w from c=0 to d=∞.
- ★ (C) Integrate in u from a=0 to b=∞ and in w from c=0 to d=u.
- (D) Integrate in u from a=0 to b=∞ and in w from c=w to d=∞.
- (E) Huh?

Convolution (6.6)

- What are the correct values for a, b, c and d?

$$\int_0^{\infty} \int_w^{\infty} e^{-su} g(w) f(u-w) du dw$$




$$= \int_a^b \int_c^d e^{-su} g(w) f(u-w) dw du$$

- (A) Integrate in u from a=0 to b=∞ and in w from c=u, d=∞.
- (B) Integrate in u from a=0 to b=w and in w from c=0 to d=∞.
- ★ (C) Integrate in u from a=0 to b=∞ and in w from c=0 to d=u.
- (D) Integrate in u from a=0 to b=∞ and in w from c=w to d=∞.
- (E) Huh?

Convolution (6.6)

- What are the correct values for a, b, c and d?

$$\int_0^{\infty} \int_w^{\infty} e^{-su} g(w) f(u-w) du dw$$




$$= \int_a^b \int_c^d e^{-su} g(w) f(u-w) dw du$$

- (A) Integrate in u from a=0 to b=∞ and in w from c=u, d=∞.
- (B) Integrate in u from a=0 to b=w and in w from c=0 to d=∞.
- ★ (C) Integrate in u from a=0 to b=∞ and in w from c=0 to d=u.
- (D) Integrate in u from a=0 to b=∞ and in w from c=w to d=∞.
- (E) Huh?

Convolution (6.6)

- What are the correct values for a, b, c and d?

$$\int_0^{\infty} \int_w^{\infty} e^{-su} g(w) f(u-w) du dw$$




$$= \int_a^b \int_c^d e^{-su} g(w) f(u-w) dw du$$

- (A) Integrate in u from a=0 to b=∞ and in w from c=u, d=∞.
- (B) Integrate in u from a=0 to b=w and in w from c=0 to d=∞.
- ★ (C) Integrate in u from a=0 to b=∞ and in w from c=0 to d=u.
- (D) Integrate in u from a=0 to b=∞ and in w from c=w to d=∞.
- (E) Huh?

Convolution (6.6)

- What are the correct values for a, b, c and d?

$$\int_0^{\infty} \int_w^{\infty} e^{-su} g(w) f(u-w) du dw$$



$$= \int_a^b \int_c^d e^{-su} g(w) f(u-w) dw du$$

- (A) Integrate in u from a=0 to b= ∞ and in w from c=u, d= ∞ .
- (B) Integrate in u from a=0 to b=w and in w from c=0 to d= ∞ .
- ★ (C) Integrate in u from a=0 to b= ∞ and in w from c=0 to d=u.
- (D) Integrate in u from a=0 to b= ∞ and in w from c=w to d= ∞ .
- (E) Huh?

Convolution (6.6)

$$\begin{aligned} F(s)G(s) &= \int_0^{\infty} e^{-s\tau} f(\tau) d\tau \int_0^{\infty} e^{-sw} g(w) dw \\ &= \int_a^b \int_c^d e^{-su} g(w) f(u-w) dw du \end{aligned}$$


Convolution (6.6)

$$\begin{aligned} F(s)G(s) &= \int_0^{\infty} e^{-s\tau} f(\tau) d\tau \int_0^{\infty} e^{-sw} g(w) dw \\ &= \int_0^{\infty} \int_0^u e^{-su} g(w) f(u-w) dw du \end{aligned}$$



Convolution (6.6)

$$\begin{aligned} F(s)G(s) &= \int_0^{\infty} e^{-s\tau} f(\tau) d\tau \int_0^{\infty} e^{-sw} g(w) dw \\ &= \int_0^{\infty} \int_0^u e^{-su} g(w) f(u-w) dw du \\ &= \int_0^{\infty} e^{-su} \int_0^u g(w) f(u-w) dw du \\ &= \int_0^{\infty} e^{-su} h(u) du = H(s) \end{aligned}$$

 where $h(u) = \int_0^u g(w) f(u-w) dw$

This is called **the convolution of f and g**.
Denoted $f * g$.

Convolution (6.6)

$$\begin{aligned} F(s)G(s) &= \int_0^{\infty} e^{-s\tau} f(\tau) d\tau \int_0^{\infty} e^{-sw} g(w) dw \\ &= \int_0^{\infty} \int_0^u e^{-su} g(w) f(u-w) dw du \\ &= \int_0^{\infty} e^{-su} \int_0^u g(w) f(u-w) dw du \\ &= \int_0^{\infty} e^{-su} h(u) du = H(s) \end{aligned}$$



The transform of a convolution is the product of the transforms.

$$h(t) = f * g(t) = \int_0^t g(w) f(t-w) dw$$

$$\Rightarrow H(s) = F(s)G(s)$$

where $h(u) = \int_0^u g(w) f(u-w) dw$

This is called **the convolution of f and g**.
Denoted $f * g$.

Convolution (6.6)

- To invert $Y(s) = \frac{1}{s^2} \cdot \frac{2}{s^2 + 4}$, we can use the fact that the inverse is the convolution of the inverses of the two pieces (instead of PFD...).

$$\mathcal{L}^{-1} \left\{ \frac{1}{s^2} \right\} =$$

$$\mathcal{L}^{-1} \left\{ \frac{2}{s^2 + 4} \right\} =$$

Convolution (6.6)

- To invert $Y(s) = \frac{1}{s^2} \cdot \frac{2}{s^2 + 4}$, we can use the fact that the inverse is the convolution of the inverses of the two pieces (instead of PFD...).

$$\mathcal{L}^{-1} \left\{ \frac{1}{s^2} \right\} = t$$

$$\mathcal{L}^{-1} \left\{ \frac{2}{s^2 + 4} \right\} =$$

Convolution (6.6)

- To invert $Y(s) = \frac{1}{s^2} \cdot \frac{2}{s^2 + 4}$, we can use the fact that the inverse is the convolution of the inverses of the two pieces (instead of PFD...).

$$\mathcal{L}^{-1} \left\{ \frac{1}{s^2} \right\} = t$$

$$\mathcal{L}^{-1} \left\{ \frac{2}{s^2 + 4} \right\} = \sin(2t)$$

Convolution (6.6)

- To invert $Y(s) = \frac{1}{s^2} \cdot \frac{2}{s^2 + 4}$, we can use the fact that the inverse is the convolution of the inverses of the two pieces (instead of PFD...).

$$\mathcal{L}^{-1} \left\{ \frac{1}{s^2} \right\} = t$$

$$\mathcal{L}^{-1} \left\{ \frac{2}{s^2 + 4} \right\} = \sin(2t)$$

$$y(t) =$$

$$(A) \int_0^t (t - w) \sin(2w) \, dw$$

$$(C) \int_0^t w \sin(2(t - w)) \, dw$$

$$(B) \int_0^t (t - w) \sin(2t) \, dw$$

$$(D) \int_0^t w \sin(2(w - t)) \, dw$$

Convolution (6.6)

- To invert $Y(s) = \frac{1}{s^2} \cdot \frac{2}{s^2 + 4}$, we can use the fact that the inverse is the convolution of the inverses of the two pieces (instead of PFD...).

$$\mathcal{L}^{-1} \left\{ \frac{1}{s^2} \right\} = t$$

$$\mathcal{L}^{-1} \left\{ \frac{2}{s^2 + 4} \right\} = \sin(2t)$$

$$y(t) =$$

$$\star (A) \int_0^t (t - w) \sin(2w) dw$$

$$\star (C) \int_0^t w \sin(2(t - w)) dw$$

$$(B) \int_0^t (t - w) \sin(2t) dw$$

$$(D) \int_0^t w \sin(2(w - t)) dw$$

Convolution (6.6)

- To invert $Y(s) = \frac{1}{s^2} \cdot \frac{2}{s^2 + 4}$, we can use the fact that the inverse is the convolution of the inverses of the two pieces (instead of PFD...).

$$\mathcal{L}^{-1} \left\{ \frac{1}{s^2} \right\} = t$$

$$\mathcal{L}^{-1} \left\{ \frac{2}{s^2 + 4} \right\} = \sin(2t)$$

$$f * g = g * f$$

$$\int_0^t f(t-w)g(w) dw = \int_0^t f(w)g(t-w) dw$$

$$y(t) =$$

$$\star (A) \int_0^t (t-w) \sin(2w) dw$$

$$\star (C) \int_0^t w \sin(2(t-w)) dw$$

$$(B) \int_0^t (t-w) \sin(2t) dw$$

$$(D) \int_0^t w \sin(2(w-t)) dw$$

Convolution (6.6)

- Transfer functions

$$ay'' + by' + cy = g(t), \quad y(0) = 0, \quad y'(0) = 0$$

Convolution (6.6)

- Transfer functions

$$ay'' + by' + cy = g(t), \quad y(0) = 0, \quad y'(0) = 0$$

$$Y(s) = \frac{1}{as^2 + bs + c}G(s)$$

Convolution (6.6)

- Transfer functions

$$ay'' + by' + cy = g(t), \quad y(0) = 0, \quad y'(0) = 0$$

$$Y(s) = \frac{1}{as^2 + bs + c}G(s)$$

- Define the transfer function for the ODE:

Convolution (6.6)

- Transfer functions

$$ay'' + by' + cy = g(t), \quad y(0) = 0, \quad y'(0) = 0$$

$$Y(s) = \frac{1}{as^2 + bs + c}G(s)$$

- Define the transfer function for the ODE:

$$H(s) = \frac{1}{as^2 + bs + c}$$

Convolution (6.6)

- Transfer functions

$$ay'' + by' + cy = g(t), \quad y(0) = 0, \quad y'(0) = 0$$

$$Y(s) = \frac{1}{as^2 + bs + c} G(s)$$

- Define the transfer function for the ODE:

$$H(s) = \frac{1}{as^2 + bs + c} \quad \text{Independent of } g(t)!$$

Convolution (6.6)

- Transfer functions

$$ay'' + by' + cy = g(t), \quad y(0) = 0, \quad y'(0) = 0$$

$$Y(s) = \frac{1}{as^2 + bs + c} G(s)$$

- Define the transfer function for the ODE:

$$H(s) = \frac{1}{as^2 + bs + c}$$

Independent of $g(t)$!

$$y(t) = (h * g)(t)$$

Convolution (6.6)

- Transfer functions

$$ay'' + by' + cy = g(t), \quad y(0) = 0, \quad y'(0) = 0$$

$$Y(s) = \frac{1}{as^2 + bs + c} G(s)$$

- Define the transfer function for the ODE:

$$H(s) = \frac{1}{as^2 + bs + c} \quad \text{Independent of } g(t)!$$

$$y(t) = (h * g)(t)$$

- $h(t)$ is called the impulse response because it solves (1) when $g(t) = \delta(t)$.

Convolution (6.6)

- Transfer functions

$$ay'' + by' + cy = g(t), \quad y(0) = 0, \quad y'(0) = 0$$

$$Y(s) = \frac{1}{as^2 + bs + c} G(s)$$

- Define the transfer function for the ODE:

$$H(s) = \frac{1}{as^2 + bs + c} \quad \text{Independent of } g(t)!$$

$$y(t) = (h * g)(t)$$

- $h(t)$ is called the impulse response because it solves (1) when $g(t) = \delta(t)$.

$$g(t) = \delta(t)$$

Convolution (6.6)

- Transfer functions

$$ay'' + by' + cy = g(t), \quad y(0) = 0, \quad y'(0) = 0$$

$$Y(s) = \frac{1}{as^2 + bs + c} G(s)$$

- Define the transfer function for the ODE:

$$H(s) = \frac{1}{as^2 + bs + c} \quad \text{Independent of } g(t)!$$

$$y(t) = (h * g)(t)$$

- $h(t)$ is called the impulse response because it solves (1) when $g(t) = \delta(t)$.

$$g(t) = \delta(t)$$

$$G(s) = e^{-0s} = 1$$

Convolution (6.6)

- Transfer functions

$$ay'' + by' + cy = g(t), \quad y(0) = 0, \quad y'(0) = 0$$

$$Y(s) = \frac{1}{as^2 + bs + c} G(s)$$

- Define the transfer function for the ODE:

$$H(s) = \frac{1}{as^2 + bs + c} \quad \text{Independent of } g(t)!$$

$$y(t) = (h * g)(t)$$

- $h(t)$ is called the impulse response because it solves (1) when $g(t) = \delta(t)$.

$$g(t) = \delta(t)$$

$$G(s) = e^{-0s} = 1$$

$$Y(s) = \frac{1}{as^2 + bs + c}$$

Convolution (6.6)

- Transfer functions

$$ay'' + by' + cy = g(t), \quad y(0) = 0, \quad y'(0) = 0$$

$$Y(s) = \frac{1}{as^2 + bs + c} G(s)$$

- Define the transfer function for the ODE:

$$H(s) = \frac{1}{as^2 + bs + c} \quad \text{Independent of } g(t)!$$

$$y(t) = (h * g)(t)$$

- $h(t)$ is called the impulse response because it solves (1) when $g(t) = \delta(t)$.

$$g(t) = \delta(t)$$

$$G(s) = \frac{e^{-0s}}{1} = 1$$

$$Y(s) = \frac{1}{as^2 + bs + c}$$

$$y_{IR}(t) = h(t) = \mathcal{L}^{-1} \left\{ \frac{1}{as^2 + bs + c} \right\}$$

Convolution (6.6)

- Transfer functions

$$ay'' + by' + cy = g(t), \quad y(0) = 0, \quad y'(0) = 0$$

$$Y(s) = \frac{1}{as^2 + bs + c} G(s)$$

- Define the transfer function for the ODE:

$$H(s) = \frac{1}{as^2 + bs + c} \quad \text{Independent of } g(t)!$$


$$y(t) = (h * g)(t)$$

- $h(t)$ is called the impulse response because it solves (1) when $g(t) = \delta(t)$.

$$g(t) = \delta(t)$$

$$G(s) = e^{-0s} = 1$$

$$Y(s) = \frac{1}{as^2 + bs + c}$$

$$y_{IR}(t) = h(t) = \mathcal{L}^{-1} \left\{ \frac{1}{as^2 + bs + c} \right\}$$


Convolution (6.6)

- Interpreting the transfer function in a model of memory.
- Your contact list got deleted. You are forced to memorize phone numbers. Let $n(t)$ be the number of phone numbers you remember at time t . You forget numbers at a rate k . Finally, $g(t)$ is the number of phone numbers per unit time that you memorize at time t .
- Equation:
- Transform of $n(t)$:
- Impulse response:

Convolution (6.6)

- Interpreting the transfer function in a model of memory.
- Your contact list got deleted. You are forced to memorize phone numbers. Let $n(t)$ be the number of phone numbers you remember at time t . You forget numbers at a rate k . Finally, $g(t)$ is the number of phone numbers per unit time that you memorize at time t .

- Equation:
$$n' = -kn + g(t)$$

- Transform of $n(t)$:
- Impulse response:

Convolution (6.6)

- Interpreting the transfer function in a model of memory.
- Your contact list got deleted. You are forced to memorize phone numbers. Let $n(t)$ be the number of phone numbers you remember at time t . You forget numbers at a rate k . Finally, $g(t)$ is the number of phone numbers per unit time that you memorize at time t .

- Equation:
$$n' = -kn + g(t)$$

- Transform of $n(t)$:
$$N(s) = \frac{G(s)}{s + k}$$

- Impulse response:

Convolution (6.6)

- Interpreting the transfer function in a model of memory.
- Your contact list got deleted. You are forced to memorize phone numbers. Let $n(t)$ be the number of phone numbers you remember at time t . You forget numbers at a rate k . Finally, $g(t)$ is the number of phone numbers per unit time that you memorize at time t .

- Equation:
$$n' = -kn + g(t)$$

- Transform of $n(t)$:
$$N(s) = \frac{G(s)}{s + k}$$

- Impulse response:
$$H(s) = \frac{1}{s + k}$$

Convolution (6.6)

- Interpreting the transfer function in a model of memory.
- Your contact list got deleted. You are forced to memorize phone numbers. Let $n(t)$ be the number of phone numbers you remember at time t . You forget numbers at a rate k . Finally, $g(t)$ is the number of phone numbers per unit time that you memorize at time t .

- Equation:
$$n' = -kn + g(t)$$

- Transform of $n(t)$:
$$N(s) = \frac{G(s)}{s + k}$$

- Impulse response:
$$H(s) = \frac{1}{s + k}$$

$$h(t) = e^{-kt}$$

Convolution (6.6)

- Interpreting the transfer function in a model of memory.
- Your contact list got deleted. You are forced to memorize phone numbers. Let $n(t)$ be the number of phone numbers you remember at time t . You forget numbers at a rate k . Finally, $g(t)$ is the number of phone numbers per unit time that you memorize at time t .

- Equation:
$$n' = -kn + g(t)$$

- Transform of $n(t)$:
$$N(s) = \frac{G(s)}{s + k}$$

- Impulse response:
$$H(s) = \frac{1}{s + k}$$

$$h(t) = e^{-kt}$$

$$n(t) = \int_0^t h(t - w)g(w) dw$$

Convolution (6.6)

- Interpreting the transfer function in a model of memory.
- Your contact list got deleted. You are forced to memorize phone numbers. Let $n(t)$ be the number of phone numbers you remember at time t . You forget numbers at a rate k . Finally, $g(t)$ is the number of phone numbers per unit time that you memorize at time t .

- Equation:
$$n' = -kn + g(t)$$

- Transform of $n(t)$:
$$N(s) = \frac{G(s)}{s + k}$$

- Impulse response:
$$H(s) = \frac{1}{s + k}$$

$$h(t) = e^{-kt}$$

$$n(t) = \int_0^t h(t-w)g(w) dw = \int_0^t e^{-k(t-w)}g(w) dw$$

Convolution (6.6)

- Interpreting the transfer function in a model of memory.
- Your contact list got deleted. You are forced to memorize phone numbers. Let $n(t)$ be the number of phone numbers you remember at time t . You forget numbers at a rate k . Finally, $g(t)$ is the number of phone numbers per unit time that you memorize at time t .
- Equation:
$$n' = -kn + g(t)$$

Convolution (6.6)

- Interpreting the transfer function in a model of memory.
- Your contact list got deleted. You are forced to memorize phone numbers. Let $n(t)$ be the number of phone numbers you remember at time t . You forget numbers at a rate k . Finally, $g(t)$ is the number of phone numbers per unit time that you memorize at time t .

- Equation:
$$n' = -kn + g(t)$$

- If you memorize one phone number at $t=0$ ($g(t)=\delta(t)$), $h(t)$ tells you what's left of that memory at time t .

$$h(t) = e^{-kt}$$

Convolution (6.6)

- Interpreting the transfer function in a model of memory.
- Your contact list got deleted. You are forced to memorize phone numbers. Let $n(t)$ be the number of phone numbers you remember at time t . You forget numbers at a rate k . Finally, $g(t)$ is the number of phone numbers per unit time that you memorize at time t .

- Equation:
$$n' = -kn + g(t)$$

- If you memorize one phone number at $t=0$ ($g(t)=\delta(t)$), $h(t)$ tells you what's left of that memory at time t .

$$h(t) = e^{-kt}$$

- If you memorize numbers over time (some complicated $g(t)$),

$$\begin{aligned} n(t) &= \int_0^t h(t-w)g(w) dw \\ &= \int_0^t e^{-(t-w)}g(w) dw \end{aligned}$$

Convolution (6.6)

- Interpreting the transfer function in a model of memory.
- Your contact list got deleted. You are forced to memorize phone numbers. Let $n(t)$ be the number of phone numbers you remember at time t . You forget numbers at a rate k . Finally, $g(t)$ is the number of phone numbers per unit time that you memorize at time t .


- Equation:
$$n' = -kn + g(t)$$

- If you memorize one phone number at $t=0$ ($g(t)=\delta(t)$), $h(t)$ tells you what's left of that memory at time t .

$$h(t) = e^{-kt}$$

- If you memorize numbers over time (some complicated $g(t)$),

$$\begin{aligned} n(t) &= \int_0^t h(t-w)g(w) dw \\ &= \int_0^t e^{-(t-w)}g(w) dw \end{aligned}$$



Convolution (6.6)

- Interpreting the transfer function in a model of memory.
- Your contact list got deleted. You are forced to memorize phone numbers. Let $n(t)$ be the number of phone numbers you remember at time t . You forget numbers at a rate k . Finally, $g(t)$ is the number of phone numbers per unit time that you memorize at time t .


- Equation:
$$n' = -kn + g(t)$$

- If you memorize one phone number at $t=0$ ($g(t)=\delta(t)$), $h(t)$ tells you what's left of that memory at time t .

$$h(t) = e^{-kt}$$

- If you memorize numbers over time (some complicated $g(t)$),

$$\begin{aligned} n(t) &= \int_0^t h(t-w)g(w) dw \\ &= \int_0^t e^{-(t-w)}g(w) dw \end{aligned}$$



Convolution (6.6)

- Interpreting the transfer function in a model of memory.
- Your contact list got deleted. You are forced to memorize phone numbers. Let $n(t)$ be the number of phone numbers you remember at time t . You forget numbers at a rate k . Finally, $g(t)$ is the number of phone numbers per unit time that you memorize at time t .


- Equation:
$$n' = -kn + g(t)$$

- If you memorize one phone number at $t=0$ ($g(t)=\delta(t)$), $h(t)$ tells you what's left of that memory at time t .

$$h(t) = e^{-kt}$$

- If you memorize numbers over time (some complicated $g(t)$),

$$\begin{aligned} n(t) &= \int_0^t h(t-w)g(w) dw \\ &= \int_0^t e^{-(t-w)}g(w) dw \end{aligned}$$



Fourier series

- Recall Method of Undetermined Coefficients for equations of the form

$$ay'' + by' + cy = f(t)$$

Fourier series

- Recall Method of Undetermined Coefficients for equations of the form

$$ay'' + by' + cy = f(t)$$


- Applicable for functions $f(t)$ that are polynomials, exponentials, sin, cos and products of those.

Fourier series

- Recall Method of Undetermined Coefficients for equations of the form

$$ay'' + by' + cy = f(t)$$

- Applicable for functions $f(t)$ that are polynomials, exponentials, sin, cos and products of those.
- How about functions like this (period but not trig)?




Fourier series

- Recall Method of Undetermined Coefficients for equations of the form

$$ay'' + by' + cy = f(t)$$

- Applicable for functions $f(t)$ that are polynomials, exponentials, sin, cos and products of those.
- How about functions like this (period but not trig)?



- What if we could construct such functions using only sine and cosine functions?