

Today

- Intro to the Laplace Transform
- Solving ODEs with forcing terms using Laplace transforms - examples

Laplace transforms - intro

- Using the Laplace Transform to solve (linear) ODEs.
- Idea:

Laplace transforms - intro

- Using the Laplace Transform to solve (linear) ODEs.

- Idea:

Unknown $y(t)$ that
satisfies some ODE




Found $y(t)$

Laplace transforms - intro

- Using the Laplace Transform to solve (linear) ODEs.


- Idea:



Laplace transforms - intro

- Using the Laplace Transform to solve (linear) ODEs.


- Idea:



Laplace transforms - intro

- Using the Laplace Transform to solve (linear) ODEs.


- Idea:



Laplace transforms - intro

- Using the Laplace Transform to solve (linear) ODEs.

- Idea:



- Laplace transform of $y(t)$: $\mathcal{L}\{y(t)\} = Y(s) = \int_0^{\infty} e^{-st} y(t) dt$

Laplace transforms - examples

- What is the Laplace transform of $y(t) = 3$?

$$\mathcal{L}\{y(t)\} = Y(s) = \int_0^{\infty} e^{-st} 3 dt$$



Laplace transforms - examples

- What is the Laplace transform of $y(t) = 3$?

$$\mathcal{L}\{y(t)\} = Y(s) = \int_0^{\infty} e^{-st} 3 dt$$



$$= -\frac{3}{s} e^{-st} \Big|_0^{\infty}$$

$$= \lim_{A \rightarrow \infty} -\frac{3}{s} e^{-st} \Big|_0^A$$

$$= -\frac{3}{s} \left(\lim_{A \rightarrow \infty} e^{-sA} - 1 \right)$$

$$= \frac{3}{s} \text{ provided } s > 0 \text{ and does not exist otherwise.}$$

Laplace transforms - examples


- What is the Laplace transform of $y(t) = 3$?

$$\begin{aligned}\mathcal{L}\{y(t)\} = Y(s) &= \int_0^{\infty} e^{-st} 3 \, dt \\ &= \frac{3}{s} \quad \text{provided } s > 0 \text{ and does not} \\ &\quad \text{exist otherwise.}\end{aligned}$$

Laplace transforms - examples

- What is the Laplace transform of $y(t) = 3$?


$$\begin{aligned}\mathcal{L}\{y(t)\} = Y(s) &= \int_0^{\infty} e^{-st} 3 \, dt \\ &= \frac{3}{s} \quad \text{provided } s > 0 \text{ and does not} \\ &\quad \text{exist otherwise.}\end{aligned}$$



Laplace transforms - examples

- What is the Laplace transform of $y(t) = 3$?




$$\begin{aligned}\mathcal{L}\{y(t)\} = Y(s) &= \int_0^{\infty} e^{-st} 3 dt \\ &= \frac{3}{s} \quad \text{provided } s > 0 \text{ and does not} \\ &\quad \text{exist otherwise.}\end{aligned}$$



Laplace transforms - examples

- What is the Laplace transform of $y(t) = 3$?

$$\begin{aligned}\mathcal{L}\{y(t)\} = Y(s) &= \int_0^{\infty} e^{-st} 3 \, dt \\ &= \frac{3}{s} \quad \text{provided } s > 0 \text{ and does not} \\ &\quad \text{exist otherwise.}\end{aligned}$$



Laplace transforms - examples

- What is the Laplace transform of $y(t) = C$?

$$\mathcal{L}\{y(t)\} = Y(s) = \int_0^{\infty} e^{-st} C dt$$

Laplace transforms - examples


- What is the Laplace transform of $y(t) = C$?

$$\begin{aligned}\mathcal{L}\{y(t)\} = Y(s) &= \int_0^{\infty} e^{-st} C \, dt \\ &= \frac{C}{s} \quad \text{provided } s > 0 \text{ and does not} \\ &\quad \text{exist otherwise.}\end{aligned}$$

Laplace transforms - examples

- What is the Laplace transform of $y(t) = C$?


$$\begin{aligned}\mathcal{L}\{y(t)\} = Y(s) &= \int_0^{\infty} e^{-st} C \, dt \\ &= \frac{C}{s} \quad \text{provided } s > 0 \text{ and does not} \\ &\quad \text{exist otherwise.}\end{aligned}$$



Laplace transforms - examples

- What is the Laplace transform of $y(t) = C$?




$$\begin{aligned}\mathcal{L}\{y(t)\} = Y(s) &= \int_0^{\infty} e^{-st} C \, dt \\ &= \frac{C}{s} \quad \text{provided } s > 0 \text{ and does not} \\ &\quad \text{exist otherwise.}\end{aligned}$$



Laplace transforms - examples

- What is the Laplace transform of $y(t) = C$?

$$\begin{aligned}\mathcal{L}\{y(t)\} = Y(s) &= \int_0^{\infty} e^{-st} C dt \\ &= \frac{C}{s} \quad \text{provided } s > 0 \text{ and does not} \\ &\quad \text{exist otherwise.}\end{aligned}$$



Laplace transforms - examples

- What is the Laplace transform of $y(t) = e^{6t}$?

$$\mathcal{L}\{y(t)\} = Y(s) = \int_0^{\infty} e^{-st} e^{6t} dt$$

Laplace transforms - examples

- What is the Laplace transform of $y(t) = e^{6t}$?

$$\mathcal{L}\{y(t)\} = Y(s) = \int_0^{\infty} e^{-st} e^{6t} dt$$

(A) $Y(s) = \frac{1}{s - 6} \quad s > 0$

(C) $Y(s) = \frac{1}{s - 6} \quad s > 6$

(B) $Y(s) = \frac{1}{6 - s} \quad s > 6$

(D) $Y(s) = \frac{1}{6 - s} \quad s > 0$

Laplace transforms - examples

- What is the Laplace transform of $y(t) = e^{6t}$?

$$\mathcal{L}\{y(t)\} = Y(s) = \int_0^{\infty} e^{-st} e^{6t} dt$$

(A) $Y(s) = \frac{1}{s-6} \quad s > 0$

★(C) $Y(s) = \frac{1}{s-6} \quad s > 6$

(B) $Y(s) = \frac{1}{6-s} \quad s > 6$

(D) $Y(s) = \frac{1}{6-s} \quad s > 0$

Laplace transforms - examples

- What is the Laplace transform of $y(t) = e^{6t}$?


$$\mathcal{L}\{y(t)\} = Y(s) = \int_0^{\infty} e^{-st} e^{6t} dt$$

(A) $Y(s) = \frac{1}{s-6} \quad s > 0$

★(C) $Y(s) = \frac{1}{s-6} \quad s > 6$

(B) $Y(s) = \frac{1}{6-s} \quad s > 6$

(D) $Y(s) = \frac{1}{6-s} \quad s > 0$



Laplace transforms - examples

- What is the Laplace transform of $y(t) = e^{6t}$?


$$\mathcal{L}\{y(t)\} = Y(s) = \int_0^{\infty} e^{-st} e^{6t} dt$$

(A) $Y(s) = \frac{1}{s-6} \quad s > 0$

★(C) $Y(s) = \frac{1}{s-6} \quad s > 6$

(B) $Y(s) = \frac{1}{6-s} \quad s > 6$

(D) $Y(s) = \frac{1}{6-s} \quad s > 0$



Laplace transforms - examples

- What is the Laplace transform of $y(t) = e^{6t}$?


$$\mathcal{L}\{y(t)\} = Y(s) = \int_0^{\infty} e^{-st} e^{6t} dt$$

(A) $Y(s) = \frac{1}{s-6} \quad s > 0$


★(C) $Y(s) = \frac{1}{s-6} \quad s > 6$

(B) $Y(s) = \frac{1}{6-s} \quad s > 6$

(D) $Y(s) = \frac{1}{6-s} \quad s > 0$



$\mathcal{L} \rightarrow$



Laplace transforms - examples

- What is the Laplace transform of $f(t) = \sin t$?

Laplace transforms - examples

- What is the Laplace transform of $f(t) = \sin t$?

$$\mathcal{L}\{f(t)\} = F(s) = \int_0^{\infty} e^{-st} \sin t \, dt$$



Laplace transforms - examples

- What is the Laplace transform of $f(t) = \sin t$?

$$\mathcal{L}\{f(t)\} = F(s) = \int_0^{\infty} e^{-st} \sin t \, dt$$

$$= e^{-st}(-\cos t) \Big|_0^{\infty} - \int_0^{\infty} (-s)e^{-st}(-\cos t) \, dt$$

$$= \lim_{A \rightarrow \infty} e^{-sA}(-\cos A) - (-1) - \int_0^{\infty} (-s)e^{-st}(-\cos t) \, dt$$

$$= 1 - s \int_0^{\infty} e^{-st} \cos t \, dt \quad s > 0$$

$$= 1 - s \left(e^{-st} \sin t \Big|_0^{\infty} + s \int_0^{\infty} e^{-st} \sin t \, dt \right)$$


$$= 1 - s^2 F(s) \quad s > 0$$

$$F(s) = \frac{1}{1 + s^2} \quad s > 0$$

$$(1 + s^2)F(s) = 1$$

Laplace transforms - examples

- What is the Laplace transform of $f(t) = \sin t$?



$$= 1 - s \left(e^{-st} \sin t \Big|_0^{\infty} + s \int_0^{\infty} e^{-st} \sin t \, dt \right)$$

$$= 1 - s^2 F(s) \quad s > 0$$

$$(1 + s^2)F(s) = 1$$

$$F(s) = \frac{1}{1 + s^2} \quad s > 0$$

Laplace transforms - examples

- What is the Laplace transform of $h(t) = \sin(\omega t)$? ($\omega > 0$)

$$\mathcal{L}\{h(t)\} = H(s) = \int_0^{\infty} e^{-st} \sin(\omega t) dt$$

- Hint: $u = \omega t$
 $du = \omega dt$

(A) $H(s) = \frac{\omega}{\omega^2 + s^2}$

(B) $H(s) = \frac{1}{1 + \left(\frac{s}{\omega}\right)^2}$

(C) $H(s) = \frac{1}{\omega} \frac{1}{1 + s^2}$

(D) $H(s) = \frac{1}{1 + s^2}$


(E) Huh?

Laplace transforms - examples

- What is the Laplace transform of $h(t) = \sin(\omega t)$? ($\omega > 0$)

$$\mathcal{L}\{h(t)\} = H(s) = \int_0^{\infty} e^{-st} \sin(\omega t) dt$$

- Hint: $u = \omega t$
 $du = \omega dt$

★ (A) $H(s) = \frac{\omega}{\omega^2 + s^2}$ 

(B) $H(s) = \frac{1}{1 + \left(\frac{s}{\omega}\right)^2}$

(C) $H(s) = \frac{1}{\omega} \frac{1}{1 + s^2}$

(D) $H(s) = \frac{1}{1 + s^2}$

(E) Huh?

Laplace transforms - examples

- What is the Laplace transform of $h(t) = \sin(\omega t)$? ($\omega > 0$)

$$\mathcal{L}\{h(t)\} = H(s) = \int_0^{\infty} e^{-st} \sin(\omega t) dt$$

• Hint: $u = \omega t$
 $du = \omega dt$

★ (A) $H(s) = \frac{\omega}{\omega^2 + s^2}$



$$H(s) = \int_0^{\infty} e^{-s \frac{u}{\omega}} \sin u \frac{du}{\omega}$$

(B) $H(s) = \frac{1}{1 + \left(\frac{s}{\omega}\right)^2}$

$$= \frac{1}{\omega} \int_0^{\infty} e^{-\frac{s}{\omega} u} \sin u du$$

(C) $H(s) = \frac{1}{\omega} \frac{1}{1 + s^2}$

$$= \frac{1}{\omega} F\left(\frac{s}{\omega}\right)$$

(D) $H(s) = \frac{1}{1 + s^2}$

(E) Huh?

$$= \frac{1}{\omega} \frac{1}{1 + \left(\frac{s}{\omega}\right)^2} \quad s > 0$$

Laplace transforms - examples

- What is the Laplace transform of $g(t) = \cos t$?
- Could calculate directly but note that $g(t) = f'(t)$ where $f(t) = \sin t$.

Laplace transforms - examples

- What is the Laplace transform of $g(t) = \cos t$?
- Could calculate directly but note that $g(t) = f'(t)$ where $f(t) = \sin t$.

$$F(s) = \mathcal{L}\{f(t)\} = \int_0^{\infty} e^{-st} f(t) dt$$

Laplace transforms - examples

- What is the Laplace transform of $g(t) = \cos t$?
- Could calculate directly but note that $g(t) = f'(t)$ where $f(t) = \sin t$.

$$F(s) = \mathcal{L}\{f(t)\} = \int_0^{\infty} e^{-st} f(t) dt$$
$$G(s) = \mathcal{L}\{g(t)\} = \int_0^{\infty} e^{-st} f'(t) dt$$



Laplace transforms - examples

- What is the Laplace transform of $g(t) = \cos t$?
- Could calculate directly but note that $g(t) = f'(t)$ where $f(t) = \sin t$.

$$F(s) = \mathcal{L}\{f(t)\} = \int_0^{\infty} e^{-st} f(t) dt$$

$$G(s) = \mathcal{L}\{g(t)\} = \int_0^{\infty} e^{-st} f'(t) dt$$



$$= e^{-st} f(t) \Big|_0^{\infty} + s \int_0^{\infty} e^{-st} f(t) dt$$

$$= -f(0) + sF(s) \quad s > 0$$

$$= -0 + s \frac{1}{1+s^2} = \frac{s}{1+s^2}$$

Laplace transforms - examples

- What is the Laplace transform of $g(t) = \cos t$?
- Could calculate directly but note that $g(t) = f'(t)$ where $f(t) = \sin t$.

$$F(s) = \mathcal{L}\{f(t)\} = \int_0^{\infty} e^{-st} f(t) dt$$

$$G(s) = \mathcal{L}\{g(t)\} = \int_0^{\infty} e^{-st} f'(t) dt$$

$$= e^{-st} f(t) \Big|_0^{\infty} + s \int_0^{\infty} e^{-st} f(t) dt$$

$$= -f(0) + sF(s) \quad s > 0$$

$$= -0 + s \frac{1}{1+s^2} = \frac{s}{1+s^2}$$

Laplace transforms - examples

- What is the Laplace transform of $\cos t$?

$$G(s) = \mathcal{L}\{\cos t\} = \frac{s}{1 + s^2}$$

- Could calculate directly but note that $g(t) = \int_0^t f(t) dt$ where $f(t) = \sin t$.

$$F(s) = \mathcal{L}\{f(t)\} = \int_0^{\infty} e^{-st} f(t) dt$$

$$G(s) = \mathcal{L}\{g(t)\} = \int_0^{\infty} e^{-st} f'(t) dt$$

$$= e^{-st} f(t) \Big|_0^{\infty} + s \int_0^{\infty} e^{-st} f(t) dt$$

$$= -f(0) + sF(s) \quad s > 0$$


$$= -0 + s \frac{1}{1 + s^2} = \frac{s}{1 + s^2}$$

Laplace transforms - examples

- What is the Laplace transform of $\cos t$?

$$G(s) = \mathcal{L}\{\cos t\} = \frac{s}{1 + s^2}$$

- Could calculate directly but note that $g(t) = \int_0^t f(t) dt$ where $f(t) = \sin t$.



$$= -f(0) + sF(s) \quad s > 0$$


$$= -0 + s \frac{1}{1 + s^2} = \frac{s}{1 + s^2}$$

Laplace transforms - examples

- What is the Laplace transform of $\cos t$?

$$G(s) = \mathcal{L}\{\cos t\} = \frac{s}{1 + s^2}$$

- Could calculate directly but note that $g(t) = f'(t)$ where $f(t) = \sin t$.



$$\mathcal{L}\{f'(t)\} = sF(s) - f(0)$$

$$\frac{s}{1 + s^2}$$

Laplace transforms - examples

- What is the Laplace transform of $h(t) = f(\omega t)$ if $\mathcal{L}\{f(t)\} = F(s)$?

(A) $H(s) = \omega F(s)$

(B) $H(s) = \frac{1}{\omega} F\left(\frac{s}{\omega}\right)$

(C) $H(s) = \omega F\left(\frac{s}{\omega}\right)$

(D) $H(s) = \frac{1}{\omega} F(s)$

(E) Don't know.

$$\mathcal{L}\{f(\omega t)\} = \int_0^{\infty} e^{-st} f(\omega t) dt$$

- Hint: $u = \omega t$
 $du = \omega dt$

Laplace transforms - examples

- What is the Laplace transform of $h(t) = f(\omega t)$ if $\mathcal{L}\{f(t)\} = F(s)$?

(A) $H(s) = \omega F(s)$

★ (B) $H(s) = \frac{1}{\omega} F\left(\frac{s}{\omega}\right)$

(C) $H(s) = \omega F\left(\frac{s}{\omega}\right)$

(D) $H(s) = \frac{1}{\omega} F(s)$

(E) Don't know.

$$\mathcal{L}\{f(\omega t)\} = \int_0^{\infty} e^{-st} f(\omega t) dt$$

- Hint: $u = \omega t$
 $du = \omega dt$

Laplace transforms - examples

- What is the Laplace transform of $h(t) = f(\omega t)$ if $\mathcal{L}\{f(t)\} = F(s)$?
- Recall two examples back:

Laplace transforms - examples

- What is the Laplace transform of $h(t) = f(\omega t)$ if $\mathcal{L}\{f(t)\} = F(s)$?
- Recall two examples back:

$$\begin{aligned}\mathcal{L}\{h(t)\} = H(s) &= \int_0^{\infty} e^{-st} \sin(\omega t) dt && u = \omega t \\ &= \int_0^{\infty} e^{-s \frac{u}{\omega}} \sin u \frac{du}{\omega} && du = \omega dt \\ &= \frac{1}{\omega} \int_0^{\infty} e^{-\frac{s}{\omega} u} \sin u du \\ &= \frac{1}{\omega} F\left(\frac{s}{\omega}\right)\end{aligned}$$

Laplace transforms - examples

- What is the Laplace transform of $h(t) = f(\omega t)$ if $\mathcal{L}\{f(t)\} = F(s)$?
- Recall two examples back:

$$\begin{aligned}\mathcal{L}\{h(t)\} = H(s) &= \int_0^{\infty} e^{-st} f(\omega t) dt && u = \omega t \\ &= \int_0^{\infty} e^{-s\frac{u}{\omega}} \sin u \frac{du}{\omega} && du = \omega dt \\ &= \frac{1}{\omega} \int_0^{\infty} e^{-\frac{s}{\omega}u} \sin u du \\ &= \frac{1}{\omega} F\left(\frac{s}{\omega}\right)\end{aligned}$$

Laplace transforms - examples

- What is the Laplace transform of $h(t) = f(\omega t)$ if $\mathcal{L}\{f(t)\} = F(s)$?
- Recall two examples back:

$$\begin{aligned}\mathcal{L}\{h(t)\} = H(s) &= \int_0^{\infty} e^{-st} f(\omega t) dt && u = \omega t \\ &= \int_0^{\infty} e^{-s\frac{u}{\omega}} f(u) \frac{du}{\omega} && du = \omega dt \\ &= \frac{1}{\omega} \int_0^{\infty} e^{-\frac{s}{\omega}u} \sin u du \\ &= \frac{1}{\omega} F\left(\frac{s}{\omega}\right)\end{aligned}$$

Laplace transforms - examples

- What is the Laplace transform of $h(t) = f(\omega t)$ if $\mathcal{L}\{f(t)\} = F(s)$?
- Recall two examples back:

$$\begin{aligned}\mathcal{L}\{h(t)\} = H(s) &= \int_0^{\infty} e^{-st} f(\omega t) dt && u = \omega t \\ &= \int_0^{\infty} e^{-s\frac{u}{\omega}} f(u) \frac{du}{\omega} && du = \omega dt \\ &= \frac{1}{\omega} \int_0^{\infty} e^{-\frac{s}{\omega}u} f(u) du \\ &= \frac{1}{\omega} F\left(\frac{s}{\omega}\right)\end{aligned}$$

Laplace transforms - examples

- What is the Laplace transform of $h(t) = f(\omega t)$ if $\mathcal{L}\{f(t)\} = F(s)$?
- Recall two examples back:

$$\begin{aligned}\mathcal{L}\{h(t)\} = H(s) &= \int_0^{\infty} e^{-st} f(\omega t) dt \\ &= \int_0^{\infty} e^{-s\frac{u}{\omega}} f(u) \frac{du}{\omega} \\ &= \frac{1}{\omega} \int_0^{\infty} e^{-\frac{s}{\omega}u} f(u) du \\ &= \frac{1}{\omega} F\left(\frac{s}{\omega}\right)\end{aligned}$$

$$\begin{aligned}u &= \omega t \\ du &= \omega dt\end{aligned}$$

$$\mathcal{L}\{\cos t\} = \frac{s}{1 + s^2}$$

$$\mathcal{L}\{\cos(\omega t)\}$$

Laplace transforms - examples

- What is the Laplace transform of $h(t) = f(\omega t)$ if $\mathcal{L}\{f(t)\} = F(s)$?
- Recall two examples back:

$$\begin{aligned}\mathcal{L}\{h(t)\} = H(s) &= \int_0^{\infty} e^{-st} f(\omega t) dt \\ &= \int_0^{\infty} e^{-s\frac{u}{\omega}} f(u) \frac{du}{\omega} \\ &= \frac{1}{\omega} \int_0^{\infty} e^{-\frac{s}{\omega}u} f(u) du \\ &= \frac{1}{\omega} F\left(\frac{s}{\omega}\right)\end{aligned}$$

$$\begin{aligned}u &= \omega t \\ du &= \omega dt\end{aligned}$$

$$\mathcal{L}\{\cos t\} = \frac{s}{1 + s^2}$$

$$\begin{aligned}\mathcal{L}\{\cos(\omega t)\} \\ &= \frac{1}{\omega} \frac{\frac{s}{\omega}}{1 + \left(\frac{s}{\omega}\right)^2}\end{aligned}$$

Laplace transforms - examples

- What is the Laplace transform of $h(t) = f(\omega t)$ if $\mathcal{L}\{f(t)\} = F(s)$?
- Recall two examples back:

$$\begin{aligned}\mathcal{L}\{h(t)\} = H(s) &= \int_0^{\infty} e^{-st} f(\omega t) dt \\ &= \int_0^{\infty} e^{-s\frac{u}{\omega}} f(u) \frac{du}{\omega} \\ &= \frac{1}{\omega} \int_0^{\infty} e^{-\frac{s}{\omega}u} f(u) du \\ &= \frac{1}{\omega} F\left(\frac{s}{\omega}\right)\end{aligned}$$

$$\begin{aligned}u &= \omega t \\ du &= \omega dt\end{aligned}$$

$$\mathcal{L}\{\cos t\} = \frac{s}{1 + s^2}$$

$$\begin{aligned}\mathcal{L}\{\cos(\omega t)\} &= \frac{1}{\omega} \frac{\frac{s}{\omega}}{1 + \left(\frac{s}{\omega}\right)^2} \\ &= \frac{s}{\omega^2 + s^2}\end{aligned}$$

Laplace transforms - examples

- What is the Laplace transform of $k(t) = e^{at} f(t)$ if $\mathcal{L}\{f(t)\} = F(s)$?

Laplace transforms - examples

- What is the Laplace transform of $k(t) = e^{at} f(t)$ if $\mathcal{L}\{f(t)\} = F(s)$?

$$\mathcal{L}\{k(t)\} = K(s) = \int_0^{\infty} e^{-st} e^{at} f(t) dt$$

Laplace transforms - examples

- What is the Laplace transform of $k(t) = e^{at} f(t)$ if $\mathcal{L}\{f(t)\} = F(s)$?

$$\begin{aligned}\mathcal{L}\{k(t)\} = K(s) &= \int_0^{\infty} e^{-st} e^{at} f(t) dt \\ &= \int_0^{\infty} e^{-(s-a)t} f(t) dt\end{aligned}$$

Laplace transforms - examples

- What is the Laplace transform of $k(t) = e^{at} f(t)$ if $\mathcal{L}\{f(t)\} = F(s)$?

$$\begin{aligned}\mathcal{L}\{k(t)\} &= K(s) = \int_0^{\infty} e^{-st} e^{at} f(t) dt \\ &= \int_0^{\infty} e^{-(s-a)t} f(t) dt \\ &= F(s - a)\end{aligned}$$

Laplace transforms - examples

- What is the Laplace transform of $k(t) = e^{at} f(t)$ if $\mathcal{L}\{f(t)\} = F(s)$?

$$\begin{aligned}\mathcal{L}\{k(t)\} &= K(s) = \int_0^{\infty} e^{-st} e^{at} f(t) dt \\ &= \int_0^{\infty} e^{-(s-a)t} f(t) dt \\ &= F(s - a)\end{aligned}$$

$$\mathcal{L}\{e^{-3t} \cos t\} =$$

$$(A) \frac{s}{1 + (s + 3)^2}$$

$$(B) \frac{1}{1 + (s + 3)^2}$$

$$(C) \frac{s + 3}{s^2 + 6s + 10}$$

$$(D) \frac{1}{s^2 + 6s + 10}$$

$\mathcal{L}\{\cos t\} = \frac{s}{1 + s^2}$

Laplace transforms - examples

- What is the Laplace transform of $k(t) = e^{at} f(t)$ if $\mathcal{L}\{f(t)\} = F(s)$?

$$\begin{aligned}\mathcal{L}\{k(t)\} = K(s) &= \int_0^{\infty} e^{-st} e^{at} f(t) dt \\ &= \int_0^{\infty} e^{-(s-a)t} f(t) dt \\ &= F(s - a)\end{aligned}$$

$\mathcal{L}\{\cos t\} = \frac{s}{1 + s^2}$

$$\mathcal{L}\{e^{-3t} \cos t\} =$$

(A) $\frac{s}{1 + (s + 3)^2}$

(B) $\frac{1}{1 + (s + 3)^2}$

★(C) $\frac{s + 3}{s^2 + 6s + 10}$

(D) $\frac{1}{s^2 + 6s + 10}$

Solving IVPs using Laplace transforms

Solving IVPs using Laplace transforms

- Solve the equation $ay'' + by' + cy = 0$ using Laplace transforms.

Solving IVPs using Laplace transforms

- Solve the equation $ay'' + by' + cy = 0$ using Laplace transforms.
- Recall that $\mathcal{L}\{f'(t)\} = sF(s) - f(0)$.

Solving IVPs using Laplace transforms

- Solve the equation $ay'' + by' + cy = 0$ using Laplace transforms.
- Recall that $\mathcal{L}\{f'(t)\} = sF(s) - f(0)$.
- Applying this to f'' , we find that



Solving IVPs using Laplace transforms

- Solve the equation $ay'' + by' + cy = 0$ using Laplace transforms.
- Recall that $\mathcal{L}\{f'(t)\} = sF(s) - f(0)$.
- Applying this to f'' , we find that

$$\begin{aligned} \pencil \mathcal{L}\{f''(t)\} &= s\mathcal{L}\{f'(t)\} - f'(0) \\ &= s(sF(s) - f(0)) - f'(0) \\ &= s^2F(s) - sf(0) - f'(0) \end{aligned}$$

Solving IVPs using Laplace transforms

- Solve the equation $ay'' + by' + cy = 0$ using Laplace transforms.
- Recall that $\mathcal{L}\{f'(t)\} = sF(s) - f(0)$.
- Applying this to f'' , we find that
$$\begin{aligned} \mathcal{L}\{f''(t)\} &= s\mathcal{L}\{f'(t)\} - f'(0) \\ &= s(sF(s) - f(0)) - f'(0) \\ &= s^2F(s) - sf(0) - f'(0) \end{aligned}$$
- Transforming both sides of the equation,

Solving IVPs using Laplace transforms

- Solve the equation $ay'' + by' + cy = 0$ using Laplace transforms.
- Recall that $\mathcal{L}\{f'(t)\} = sF(s) - f(0)$.

- Applying this to f'' , we find that

$$\begin{aligned} \pencil \mathcal{L}\{f''(t)\} &= s\mathcal{L}\{f'(t)\} - f'(0) \\ &= s(sF(s) - f(0)) - f'(0) \\ &= s^2F(s) - sf(0) - f'(0) \end{aligned}$$

- Transforming both sides of the equation,

$$\mathcal{L}\{ay'' + by' + cy\} = 0$$



Solving IVPs using Laplace transforms

- Solve the equation $ay'' + by' + cy = 0$ using Laplace transforms.

- Recall that $\mathcal{L}\{f'(t)\} = sF(s) - f(0)$.

- Applying this to f'' , we find that


$$\begin{aligned} \pencil \mathcal{L}\{f''(t)\} &= s\mathcal{L}\{f'(t)\} - f'(0) \\ &= s(sF(s) - f(0)) - f'(0) \\ &= s^2F(s) - sf(0) - f'(0) \end{aligned}$$

- Transforming both sides of the equation,

$$\mathcal{L}\{ay'' + by' + cy\} = 0 \qquad Y(s) = \frac{asy(0) + ay'(0) + by(0)}{as^2 + bs + c}$$

$$\pencil \underline{a\mathcal{L}\{y''\}} + \underline{b\mathcal{L}\{y'\}} + \underline{c\mathcal{L}\{y\}} = 0$$

$$a(\underline{s^2Y(s) - sy(0) - y'(0)}) + b(\underline{sY(s) - y(0)}) + \underline{cY(s)} = 0$$

$$(as^2 + bs + c)Y(s) = asy(0) + ay'(0) + by(0)$$


Solving IVPs using Laplace transforms

- Solve the equation $y'' + 4y = 0$ with initial conditions $y(0)=1$, $y'(0)=0$ using Laplace transforms.

Solving IVPs using Laplace transforms

- Solve the equation $y'' + 4y = 0$ with initial conditions $y(0)=1$, $y'(0)=0$ using Laplace transforms.

$$s^2 Y(s) - sy(0) - y'(0) + 4Y(s) = 0$$

$$s^2 Y(s) - s - 0 + 4Y(s) = 0$$

$$s^2 Y(s) + 4Y(s) = s$$

$$Y(s) = \frac{s}{s^2 + 4}$$

Solving IVPs using Laplace transforms

- Solve the equation $y'' + 4y = 0$ with initial conditions $y(0)=1$, $y'(0)=0$ using Laplace transforms.

$$s^2 Y(s) - sy(0) - y'(0) + 4Y(s) = 0$$

$$s^2 Y(s) - s - 0 + 4Y(s) = 0$$

$$s^2 Y(s) + 4Y(s) = s$$

$$Y(s) = \frac{s}{s^2 + 4}$$

- To find $y(t)$, we have to invert the transform. What $y(t)$ would have $Y(s)$ as its transform?

Solving IVPs using Laplace transforms

- Solve the equation $y'' + 4y = 0$ with initial conditions $y(0)=1$, $y'(0)=0$ using Laplace transforms.

$$s^2 Y(s) - sy(0) - y'(0) + 4Y(s) = 0$$

$$s^2 Y(s) - s - 0 + 4Y(s) = 0$$

$$s^2 Y(s) + 4Y(s) = s$$

$$Y(s) = \frac{s}{s^2 + 4}$$

- To find $y(t)$, we have to invert the transform. What $y(t)$ would have $Y(s)$ as its transform?

- Recall that $\mathcal{L}\{\cos(\omega t)\} = \frac{s}{\omega^2 + s^2}$. So $y(t) = \cos(2t)$.

Solving IVPs using Laplace transforms

- Solve the equation $y'' + 6y' + 13y = 0$ with initial conditions $y(0)=1$, $y'(0)=0$ using Laplace transforms.



Solving IVPs using Laplace transforms

- Solve the equation $y'' + 6y' + 13y = 0$ with initial conditions $y(0)=1$, $y'(0)=0$ using Laplace transforms.



$$Y(s) = \frac{s + 6}{s^2 + 6s + 13}$$

Solving IVPs using Laplace transforms

- Solve the equation $y'' + 6y' + 13y = 0$ with initial conditions $y(0)=1$, $y'(0)=0$ using Laplace transforms.



$$Y(s) = \frac{s + 6}{s^2 + 6s + 13}$$

- To find $y(t)$, we have to invert the transform. What $y(t)$ would have $Y(s)$ as its transform?

Solving IVPs using Laplace transforms

- Solve the equation $y'' + 6y' + 13y = 0$ with initial conditions $y(0)=1$, $y'(0)=0$ using Laplace transforms.



$$Y(s) = \frac{s + 6}{s^2 + 6s + 13}$$

- To find $y(t)$, we have to invert the transform. What $y(t)$ would have $Y(s)$ as its transform?

$$\mathcal{L}\{\cos(\omega t)\} = \frac{s}{s^2 + \omega^2}$$

$$\mathcal{L}\{\sin(\omega t)\} = \frac{\omega}{s^2 + \omega^2}$$

$$\mathcal{L}\{e^{at} f(t)\} = F(s - a)$$

$$\mathcal{L}\{e^{-3t} \cos t\} = \frac{s + 3}{1 + (s + 3)^2}$$

Solving IVPs using Laplace transforms

- Solve the equation $y'' + 6y' + 13y = 0$ with initial conditions $y(0)=1$, $y'(0)=0$ using Laplace transforms.



$$Y(s) = \frac{s + 6}{s^2 + 6s + 13}$$

- To find $y(t)$, we have to invert the transform. What $y(t)$ would have $Y(s)$ as its transform?

$$\mathcal{L}\{\cos(\omega t)\} = \frac{s}{s^2 + \omega^2}$$

$$\mathcal{L}\{\sin(\omega t)\} = \frac{\omega}{s^2 + \omega^2}$$



$$\mathcal{L}\{e^{at} f(t)\} = F(s - a)$$

$$\mathcal{L}\{e^{-3t} \cos t\} = \frac{s + 3}{1 + (s + 3)^2}$$

Solving IVPs using Laplace transforms

- Solve the equation $y'' + 6y' + 13y = 0$ with initial conditions $y(0)=1$, $y'(0)=0$ using Laplace transforms.



$$Y(s) = \frac{s + 6}{s^2 + 6s + 13}$$

- To find $y(t)$, we have to invert the transform. What $y(t)$ would have $Y(s)$ as its transform?

$$\mathcal{L}\{\cos(\omega t)\} = \frac{s}{s^2 + \omega^2}$$

$$\mathcal{L}\{\sin(\omega t)\} = \frac{\omega}{s^2 + \omega^2}$$

$$\mathcal{L}\{e^{at} f(t)\} = F(s - a)$$

$$\mathcal{L}\{e^{-3t} \cos t\} = \frac{s + 3}{1 + (s + 3)^2}$$

$$Y(s) = \frac{s + 3 + 3}{s^2 + 6s + 9 + 4}$$

$$= \frac{s + 3}{(s + 3)^2 + 4} + \frac{3}{(s + 3)^2 + 4}$$

$$= \frac{s + 3}{(s + 3)^2 + 4} + \frac{3}{2} \frac{2}{(s + 3)^2 + 4}$$

$$y(t) = e^{-3t} \cos(2t) + \frac{3}{2} e^{-3t} \sin(2t)$$

Solving IVPs using Laplace transforms

- Solve the equation $y'' + 6y' + 13y = 0$ with initial conditions $y(0)=1$, $y'(0)=0$ using Laplace transforms.



$$Y(s) = \frac{s + 6}{s^2 + 6s + 13}$$

- To find $y(t)$, we have $\lambda = \frac{-6 \pm i\sqrt{52 - 36}}{2} = -3 \pm 2i$ would have $Y(s)$ as its transform?

$$\mathcal{L}\{\cos(\omega t)\} = \frac{s}{s^2 + \omega^2}$$

$$\mathcal{L}\{\sin(\omega t)\} = \frac{\omega}{s^2 + \omega^2}$$

$$\mathcal{L}\{e^{at} f(t)\} = F(s - a)$$

$$\mathcal{L}\{e^{-3t} \cos t\} = \frac{s + 3}{1 + (s + 3)^2}$$



$$Y(s) = \frac{s + 3}{s^2 + 6s + 9 + 4}$$

$$= \frac{s + 3}{(s + 3)^2 + 4} + \frac{3}{(s + 3)^2 + 4}$$

$$= \frac{s + 3}{(s + 3)^2 + 4} + \frac{3}{2} \frac{2}{(s + 3)^2 + 4}$$

$$y(t) = e^{-3t} \cos(2t) + \frac{3}{2} e^{-3t} \sin(2t)$$

Solving IVPs using Laplace transforms

- Solve the equation $y'' + 6y' + 13y = 0$ with initial conditions $y(0)=1$, $y'(0)=0$ using Laplace transforms.

$$Y(s) = \frac{s + 6}{s^2 + 6s + 13}$$

Solving IVPs using Laplace transforms

- Solve the equation $y'' + 6y' + 13y = 0$ with initial conditions $y(0)=1$, $y'(0)=0$ using Laplace transforms.

$$Y(s) = \frac{s + 6}{s^2 + 6s + 13}$$

1. Does the denominator have real or complex roots?

Solving IVPs using Laplace transforms

- Solve the equation $y'' + 6y' + 13y = 0$ with initial conditions $y(0)=1$, $y'(0)=0$ using Laplace transforms.

$$Y(s) = \frac{s + 6}{s^2 + 6s + 13}$$

1. Does the denominator have real or complex roots? Complex.

Solving IVPs using Laplace transforms

- Solve the equation $y'' + 6y' + 13y = 0$ with initial conditions $y(0)=1$, $y'(0)=0$ using Laplace transforms.

$$Y(s) = \frac{s + 6}{s^2 + 6s + 13} = \frac{s + 6}{s^2 + 6s + 9 + 4}$$

1. Does the denominator have real or complex roots? Complex.
2. Complete the square.

Solving IVPs using Laplace transforms

- Solve the equation $y'' + 6y' + 13y = 0$ with initial conditions $y(0)=1$, $y'(0)=0$ using Laplace transforms.

$$Y(s) = \frac{s + 6}{s^2 + 6s + 13} = \frac{s + 6}{s^2 + 6s + 9 + 4} = \frac{s + 6}{(s + 3)^2 + 4}$$

1. Does the denominator have real or complex roots? Complex.
2. Complete the square.

Solving IVPs using Laplace transforms

- Solve the equation $y'' + 6y' + 13y = 0$ with initial conditions $y(0)=1$, $y'(0)=0$ using Laplace transforms.

$$Y(s) = \frac{s+6}{s^2+6s+13} = \frac{s+6}{s^2+6s+9+4} = \frac{s+6}{(s+3)^2+4} = \frac{s+3+3}{(s+3)^2+4}$$

1. Does the denominator have real or complex roots? Complex.
2. Complete the square.
3. Put numerator in form $(s+\alpha)+\beta$ where $(s+\alpha)$ is the completed square.

Solving IVPs using Laplace transforms

- Solve the equation $y'' + 6y' + 13y = 0$ with initial conditions $y(0)=1$, $y'(0)=0$ using Laplace transforms.

$$\begin{aligned} Y(s) &= \frac{s+6}{s^2+6s+13} = \frac{s+6}{s^2+6s+9+4} = \frac{s+6}{(s+3)^2+4} = \frac{s+3+3}{(s+3)^2+4} \\ &= \frac{s+3}{(s+3)^2+4} + \frac{3}{(s+3)^2+4} \end{aligned}$$

1. Does the denominator have real or complex roots? Complex.
2. Complete the square.
3. Put numerator in form $(s+\alpha)+\beta$ where $(s+\alpha)$ is the completed square.

Solving IVPs using Laplace transforms

- Solve the equation $y'' + 6y' + 13y = 0$ with initial conditions $y(0)=1$, $y'(0)=0$ using Laplace transforms.

$$\begin{aligned} Y(s) &= \frac{s+6}{s^2+6s+13} = \frac{s+6}{s^2+6s+9+4} = \frac{s+6}{(s+3)^2+4} = \frac{s+3+3}{(s+3)^2+4} \\ &= \frac{s+3}{(s+3)^2+4} + \frac{3}{(s+3)^2+4} = \frac{s+3}{(s+3)^2+2^2} + \frac{3}{2} \frac{2}{(s+3)^2+2^2} \end{aligned}$$

1. Does the denominator have real or complex roots? Complex.
2. Complete the square.
3. Put numerator in form $(s+\alpha)+\beta$ where $(s+\alpha)$ is the completed square.
4. Fix up coefficient of the term with no s in the numerator.

Solving IVPs using Laplace transforms

- Solve the equation $y'' + 6y' + 13y = 0$ with initial conditions $y(0)=1$, $y'(0)=0$ using Laplace transforms.

$$\begin{aligned} Y(s) &= \frac{s+6}{s^2+6s+13} = \frac{s+6}{s^2+6s+9+4} = \frac{s+6}{(s+3)^2+4} = \frac{s+3+3}{(s+3)^2+4} \\ &= \frac{s+3}{(s+3)^2+4} + \frac{3}{(s+3)^2+4} = \frac{s+3}{(s+3)^2+2^2} + \frac{3}{2} \frac{2}{(s+3)^2+2^2} \end{aligned}$$

$$y(t) = e^{-3t} \cos(2t) + \frac{3}{2} e^{-3t} \sin(2t)$$

1. Does the denominator have real or complex roots? Complex.
2. Complete the square.
3. Put numerator in form $(s+\alpha)+\beta$ where $(s+\alpha)$ is the completed square.
4. Fix up coefficient of the term with no s in the numerator.
5. Invert.