

Today


- Forced vibrations
 - Newton's 2nd Law with external forcing.
 - Forced mass-spring system without damping away from resonance.
 - Forced mass-spring system without damping at resonance.
 - Forced mass-spring system with damping.
- Midterm (Feb 10, in class) - Everything up to and including Monday Feb 3 (systems of equations and review of eigenvectors).

Forced vibrations (3.8)

- Newton's 2nd Law:

$$ma = -kx - \gamma v + F(t)$$

spring force drag force applied/external force



$$mx'' + \gamma x' + kx = F(t)$$

- Forced vibrations - nonhomogeneous linear equation with constant coefficients.
- Building during earthquake, tuning fork near instrument, car over washboard road, polar bond under EM stimulus (classical, not quantum).

Forced vibrations (3.8)

- Without damping ($\gamma = 0$).

$$mx'' + kx = F_0 \cos(\omega t)$$

forcing frequency



- For what value(s) of w does this equation have an unbounded solution?

(A) $w = \text{sqrt}(k/m)$

(B) $w = m/F_0$

(C) $w = (k/m)^2$

(D) $w = 2\pi$

Forced vibrations (3.8)

- Without damping ($\gamma = 0$).

$$mx'' + kx = F_0 \cos(\omega t)$$

forcing frequency

$$mx'' + kx = 0$$

$$x_h(t) = C_1 \cos(\omega_0 t) + C_2 \sin(\omega_0 t)$$

$$\omega_0 = \sqrt{\frac{k}{m}}$$

- Case 1: $\omega \neq \omega_0$

$$x_p(t) = A \cos(\omega t) + B \sin(\omega t)$$

$$A = ?, B = ?$$

natural frequency

Forced vibrations (3.8)

- Without damping ($\gamma = 0$).

$$mx'' + kx = F_0 \cos(\omega t)$$

forcing frequency

$$mx'' + kx = 0$$

$$x_h(t) = C_1 \cos(\omega_0 t) + C_2 \sin(\omega_0 t)$$

$$\omega_0 = \sqrt{\frac{k}{m}}$$

- Case 1: $\omega \neq \omega_0$

$$x_p(t) = A \cos(\omega t) + B \sin(\omega t)$$

$$x_p''(t) = -\omega^2 A \cos(\omega t) - \omega^2 B \sin(\omega t)$$


$$mx_p'' + kx_p = (k - \omega^2 m)A \cos(\omega t) + (k - \omega^2 m)B \sin(\omega t)$$

$$= F_0 \cos(\omega t) \Rightarrow A = \frac{F_0}{k(\omega_0^2 - \omega^2)}, B = 0$$

natural frequency

Forced vibrations (3.8)

- Without damping ($\gamma = 0$), $\omega \neq \omega_0$.
 - Beats - long term behaviour includes both x_h and x_p
 - On the board.



Forced vibrations (3.8)

- Case 2: $\omega = \omega_0$

$$mx'' + kx = F_0 \cos(\omega_0 t) \quad \omega_0 = \sqrt{\frac{k}{m}}$$

~~$$x_p(t) = t(A \cos(\omega_0 t) + B \sin(\omega_0 t))$$~~

$$x'_p(t) = A \cos(\omega_0 t) + B \sin(\omega_0 t)$$

$$+ t(-\omega_0 A \sin(\omega_0 t) + \omega_0 B \cos(\omega_0 t))$$

$$x''_p(t) = -\omega_0 A \sin(\omega_0 t) + \omega_0 B \cos(\omega_0 t)$$

$$+ (-\omega_0 A \sin(\omega_0 t) + \omega_0 B \cos(\omega_0 t))$$

~~$$+ t(-\omega_0^2 A \cos(\omega_0 t) - \omega_0^2 B \sin(\omega_0 t))$$~~


$$A = 0$$

$$B = \frac{F_0}{2\omega_0 m} = \frac{F_0}{2\sqrt{km}}$$

$$x_p(t) = \frac{F_0}{2\sqrt{km}} t \sin(\omega_0 t)$$


Forced vibrations (3.8)

- Without damping ($\gamma = 0$), $\omega \neq \omega_0$.
 - Long term behaviour - x_p grows unbounded, swamping out x_h .



Forced vibrations (3.8)

- Plot of the amplitude of the particular solution as a function of ω .



- Calculated:

$$A = \frac{F_0}{m(\omega_0^2 - \omega^2)}$$

- Plotted with:

$$\frac{F_0}{m} = 1, \quad \omega_0 = 1$$

$$A(\omega) = \frac{1}{|\omega_0^2 - \omega^2|}$$

- Recall that for $\omega = \omega_0$, the amplitude grows without bound.

Forced vibrations (3.8)

- With damping (on the blackboard)
- Desmos illustration