

Today

- Introduction to systems of equations
- Direction fields
- Eigenvalues and eigenvectors
- Finding the general solution (distinct e-value case)
- Pre-midterm office hours poll - Friday (best time other than 2-3),
Monday (holiday so buildings locked)

Introduction to systems of equations

- So far, we've only dealt with equations with one unknown function. Sometimes, we'll be interested in more than one unknown function.
- Examples:
 - position of object in one dimensional space in terms of x , v :

$$mx'' + \gamma x' + kx = 0 \rightarrow mv' + \gamma v + kx = 0$$

$$x' = v \qquad v' = -\frac{\gamma}{m}v - \frac{k}{m}x$$

$$x'' = v'$$

$$x' = v$$

$$v' = -\frac{k}{m}x - \frac{\gamma}{m}v$$

$$\begin{pmatrix} x \\ v \end{pmatrix}' = \begin{pmatrix} 0 & 1 \\ -\frac{k}{m} & -\frac{\gamma}{m} \end{pmatrix} \begin{pmatrix} x \\ v \end{pmatrix}$$



Introduction to systems of equations

- So far, we've only dealt with equations with one unknown function. Sometimes, we'll be interested in more than one unknown function.
- Examples:
 - position of object in one dimensional space in terms of x , v .
 - position of an object in a plane (x , y coordinates) or three dimensional space (x , y , z coordinates).
 - positions of multiple objects (two or more masses linked by springs).
 - concentration in connected chambers (saltwater in multiple tanks, intracellular and extracellular, blood stream and organs).
 - populations of two species (e.g. predator and prey).

Introduction to systems of equations

- As with single equations, we have **linear** and **nonlinear** systems:

$$\begin{aligned}\frac{dx}{dt} &= t^2 x - y + \cos(2t) \\ \frac{dy}{dt} &= x + 4 \sin(t)y + t^3\end{aligned}$$

$$\begin{aligned}\frac{dx}{dt} &= t^2 x - y^2 \\ \frac{dy}{dt} &= \sqrt{x} - y\end{aligned}$$

- And we also have **nonhomogeneous** and **homogeneous** systems.

$$\begin{aligned}\frac{dx}{dt} &= t^2 x - y + \cos(2t) \\ \frac{dy}{dt} &= x + 4 \sin(t)y + t^3\end{aligned}$$

$$\begin{aligned}\frac{dx}{dt} &= t^2 x - y \\ \frac{dy}{dt} &= x + 4 \sin(t)y\end{aligned}$$

Introduction to systems of equations

- Any linear system can be written in matrix form:

$$\frac{dx}{dt} = t^2 x - y + \cos(2t)$$

$$\frac{dy}{dt} = x + 4 \sin(t)y + t^3$$

$$\frac{d}{dt} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} t^2 & -1 \\ 1 & 4 \sin(t) \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} \cos(2t) \\ t^3 \end{pmatrix}$$

- We'll focus on the case in which the matrix has constant entries. And homogeneous, to start. For example,

$$\frac{d}{dt} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 4 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$$

Introduction to systems of equations


- Geometric interpretation - **direction fields**.

$$\frac{d}{dt} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 4 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \quad \text{or} \quad \mathbf{x}' = A\mathbf{x}$$

- Think of the unknown functions as coordinates $(x(t), y(t))$ of an object in the plane.
- $A\mathbf{x}$ gives the velocity vector of the object located at \mathbf{x} .

$$\mathbf{x} = \begin{pmatrix} 2 \\ 1 \end{pmatrix} \quad \rightarrow$$
$$A\mathbf{x} = \begin{pmatrix} 1 & 1 \\ 4 & 1 \end{pmatrix} \begin{pmatrix} 2 \\ 1 \end{pmatrix} = \begin{pmatrix} 3 \\ 9 \end{pmatrix}$$

- Solutions must follow the arrows.



Introduction to systems of equations

- Which of the following equations matches the given direction field?


(A) $\mathbf{x}' = \begin{pmatrix} -1 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$

(B) $\mathbf{x}' = \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$

(C) $\mathbf{x}' = \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$

★ (D) $\mathbf{x}' = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$

(E) Explain, please.




[http://kevinmehall.net/p/equationexplorer/vectorfield.html#\(x+y\)i+\(x-y\)j%7C%5B-10,10,-10,10%5D](http://kevinmehall.net/p/equationexplorer/vectorfield.html#(x+y)i+(x-y)j%7C%5B-10,10,-10,10%5D)

Introduction to systems of equations

- You should see two “special” directions.
- What are they?
- Directions along which the velocity vector is parallel to the position vector.
- That is, $A\mathbf{v} = \lambda\mathbf{v}$.

$$\lambda_2 = \sqrt{2}/2$$

$$\mathbf{v}_2 = \begin{pmatrix} 1 & -1\sqrt{2} \\ \sqrt{2} & 1-1 \end{pmatrix}$$



Matrix review (eigen-calculations)

- Find eigenvalues and eigenvectors of $A = \begin{pmatrix} 1 & 1 \\ 4 & 1 \end{pmatrix}$.
- Looking for values λ and vectors \mathbf{v} for which $A\mathbf{v} = \lambda\mathbf{v}$.
- What are the eigenvalues of A ?

(A) 1 and -3

★ (B) -1 and 3

(C) 1 and 3

(D) -1 and -3

(E) Explain, please.

Matrix review (eigen-calculations)

- Find eigenvalues and eigenvectors of $A = \begin{pmatrix} 1 & 1 \\ 4 & 1 \end{pmatrix}$.
- Looking for values λ and vectors \mathbf{v} for which $A\mathbf{v} = \lambda\mathbf{v}$.


$$A\mathbf{v} - \lambda\mathbf{v} = \mathbf{0}$$

$$(A - \lambda I)\mathbf{v} = \mathbf{0}$$

$$\det(A - \lambda I) = 0$$

$$\det \begin{pmatrix} 1 - \lambda & 1 \\ 4 & 1 - \lambda \end{pmatrix} = 0$$

$$(1 - \lambda)^2 - 4 = 0$$

$$(\lambda^2 - 2\lambda - 3 = 0)$$

$$\lambda = 1 \pm 2 = -1, 3$$

- What are the eigenvectors associated with $\lambda_1 = -1$?

$$(A) \mathbf{v}_1 = \begin{pmatrix} 1 \\ -2 \end{pmatrix}$$

$$\star (B) \mathbf{v}_1 = c \begin{pmatrix} 1 \\ -2 \end{pmatrix}$$

$$(C) \mathbf{v}_1 = \begin{pmatrix} 2 \\ 1 \end{pmatrix}$$

$$(D) \mathbf{v}_1 = c \begin{pmatrix} 2 \\ 1 \end{pmatrix}$$

(E) Explain, please.

Matrix review (eigen-calculations)

- Find eigenvalues and eigenvectors of $A = \begin{pmatrix} 1 & 1 \\ 4 & 1 \end{pmatrix}$.
- Looking for values λ and vectors \mathbf{v} for which $A\mathbf{v} = \lambda\mathbf{v}$.

$$A\mathbf{v} - \lambda\mathbf{v} = \mathbf{0}$$

$$\pencil \lambda_1 = -1$$

$$(A - \lambda I)\mathbf{v} = \mathbf{0}$$

$$(A + I)\mathbf{v}_1 = \begin{pmatrix} 2 & 1 \\ 4 & 2 \end{pmatrix} \mathbf{v}_1 = \mathbf{0}$$

$$\det(A - \lambda I) = 0$$

$$\det \begin{pmatrix} 1 - \lambda & 1 \\ 4 & 1 - \lambda \end{pmatrix} = 0$$

$$\begin{pmatrix} 2 & 1 \\ 4 & 2 \end{pmatrix} \sim \begin{pmatrix} 2 & 1 \\ 0 & 0 \end{pmatrix}$$

$$(1 - \lambda)^2 - 4 = 0$$

$$2v_1 + v_2 = 0$$

$$(\lambda^2 - 2\lambda - 3 = 0)$$

$$\lambda = 1 \pm 2 = -1, 3$$

$$\mathbf{v}_1 = \begin{pmatrix} 1 \\ -2 \end{pmatrix}$$

(and any scalar multiple of it)

Matrix review (eigen-calculations)

- Find eigenvalues and eigenvectors of $A = \begin{pmatrix} 1 & 1 \\ 4 & 1 \end{pmatrix}$.
- Looking for values λ and vectors \mathbf{v} for which $A\mathbf{v} = \lambda\mathbf{v}$.

$$A\mathbf{v} - \lambda\mathbf{v} = \mathbf{0}$$

$$(A - \lambda I)\mathbf{v} = \mathbf{0}$$

$$\det(A - \lambda I) = 0$$

$$\det \begin{pmatrix} 1 - \lambda & 1 \\ 4 & 1 - \lambda \end{pmatrix} = 0$$

$$(1 - \lambda)^2 - 4 = 0$$

$$(\lambda^2 - 2\lambda - 3 = 0)$$

$$\lambda = 1 \pm 2 = -1, 3$$

$$\lambda_1 = -1$$

$$\mathbf{v}_1 = \begin{pmatrix} 1 \\ -2 \end{pmatrix}$$

$$\lambda_2 = 3$$

$$\mathbf{v}_2 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

- How do we use eigenvalues and eigenvectors to construct a general solution?

Solving a system of ODEs

- The following is a shortcut approach for 2x2 systems, mostly for insight.
- Find the general solution to the system of equations

$$\begin{aligned}x_1' &= x_1 + x_2 \\x_2' &= 4x_1 + x_2\end{aligned}\quad \text{or equivalently} \quad \mathbf{x}' = \begin{pmatrix} 1 & 1 \\ 4 & 1 \end{pmatrix} \mathbf{x}$$

- Convert this into a second order equation in only one unknown (x_1):

$$\pencil x_1'' = x_1' + x_2' = x_1' + 4x_1 + x_2$$

$$x_2 = x_1' - x_1$$

$$x_1'' = x_1' + 4x_1 + x_1' - x_1$$

$$x_1'' - 2x_1' - 3x_1 = 0$$

Solving a system of ODEs

- The following is a shortcut approach for 2x2 systems, mostly for insight.
- Find the general solution to the system of equations

$$\begin{aligned}x_1' &= x_1 + x_2 \\x_2' &= 4x_1 + x_2\end{aligned}\quad \text{or equivalently} \quad \mathbf{x}' = \begin{pmatrix} 1 & 1 \\ 4 & 1 \end{pmatrix} \mathbf{x}$$

- Convert this into a second order equation in only one unknown (x_1):

$$x_1'' - 2x_1' - 3x_1 = 0 \quad \rightarrow \quad r^2 - 2r - 3 = 0$$

$$r = -1, 3$$

$$x_1 = C_1 e^{-t} + C_2 e^{3t}$$

 $x_2 = x_1' - x_1 = -C_1 e^{-t} + 3C_2 e^{3t} - C_1 e^{-t} - C_2 e^{3t}$
 $= -2C_1 e^{-t} + 2C_2 e^{3t}$

Solving a system of ODEs

- The following is a shortcut approach for 2x2 systems, mostly for insight.
- Find the general solution to the system of equations

$$\begin{aligned}x_1' &= x_1 + x_2 \\x_2' &= 4x_1 + x_2\end{aligned}\quad \text{or equivalently} \quad \mathbf{x}' = \begin{pmatrix} 1 & 1 \\ 4 & 1 \end{pmatrix} \mathbf{x}$$

- Convert this into a second order equation in only one unknown (x_1):

$$x_1'' - 2x_1' - 3x_1 = 0 \quad \rightarrow \quad r^2 - 2r - 3 = 0$$

$$r = -1, 3$$

$$x_1 = C_1 e^{-t} + C_2 e^{3t}$$

$$x_2 = -2C_1 e^{-t} + 2C_2 e^{3t}$$

$$\mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = C_1 e^{-t} \begin{pmatrix} 1 \\ -2 \end{pmatrix} + C_2 e^{3t} \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

- Recall:

$$\lambda_1 = -1$$
$$\mathbf{v}_1 = \begin{pmatrix} 1 \\ -2 \end{pmatrix}$$

$$\lambda_2 = 3$$
$$\mathbf{v}_2 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

Solving a system of ODEs

- You can use the second order trick for 2x2 but in general,
 - Find eigenvalues and eigenvectors of A,
 - Assemble general solution by summing up terms of the form

$$C_n e^{\lambda_n t} \mathbf{v}_n$$

- This works when eigenvalues are distinct or, if there are repeated eigenvalues, when there are N independent eigenvectors.
- Other cases (not enough e-vectors or complex e-values) Thursday.